

**CALIFORNIA DEPARTMENT OF MENTAL HEALTH
OFFICE OF SUICIDE PREVENTION
DATA SUMMARY SHEET ON SUICIDE DEATHS AND NONFATAL SELF-INFLICTED INJURIES
STANISLAUS COUNTY**


YEAR 2009

CALIFORNIA DATA

| | |
|-------------------------------------|------------|
| POPULATION | 38,688,293 |
| SUICIDE DEATHS | 3,760 |
| SUICIDE DEATH RATE | 9.7 |
| NONFATAL SELF-INFLICTED INJURIES | 16,554 |
| NONFATAL SELF-INFLICTED INJURY RATE | 42.8 |

COUNTY: Stanislaus

| | |
|-------------------------------------|---------|
| POPULATION | 549,408 |
| SUICIDE DEATHS | 58 |
| SUICIDE DEATH RATE | 10.6 |
| NONFATAL SELF-INFLICTED INJURIES | 244 |
| NONFATAL SELF-INFLICTED INJURY RATE | 44.4 |



**ADDITIONAL COUNTY DATA
SUICIDE DEATHS BY AGE, GENDER & RACE/ETHNICITY**

| AGE GROUPS | 5-14 | 15-24 | 25-34 | 35-44 | 45-54 | 55-64 | 65-74 | 75-84 | 85+ |
|--------------------|--------|--------|--------|--------|--------|--------|--------|--------|-------|
| POPULATION | 97,534 | 91,826 | 71,469 | 65,267 | 69,943 | 51,301 | 29,359 | 18,705 | 7,711 |
| SUICIDE DEATHS | 0 | 5 | 8 | 16 | 15 | 8 | 2 | 3 | 1 |
| SUICIDE DEATH RATE | 0.0 | 5.4 | 11.2 | 24.5 | 21.4 | 15.6 | 6.8 | 16.0 | 13.0 |

| GENDER | FEMALES | MALES |
|--------------------|---------|---------|
| POPULATION | 279,687 | 269,721 |
| SUICIDE DEATHS | 12 | 46 |
| SUICIDE DEATH RATE | 4.3 | 17.1 |

| RACE/ETHNICITY | AMERICAN INDIAN / ALASKAN NATIVE | ASIAN | BLACK | HISPANIC | HAWAIIAN / PACIFIC ISLANDER | WHITE | TWO OR MORE RACES |
|--------------------|----------------------------------|--------|--------|----------|-----------------------------|---------|-------------------|
| POPULATION | 4,568 | 26,458 | 13,556 | 226,193 | 1,844 | 266,480 | 10,309 |
| SUICIDE DEATHS | 0 | 0 | 1 | 10 | 1 | 44 | 2 |
| SUICIDE DEATH RATE | 0.0 | 0.0 | 7.4 | 4.4 | 54.2 | 16.5 | 19.4 |

* 1) Use caution when analyzing small numbers and rates derived from them; 2) Rates calculated from small numbers of deaths or from small populations tend to be **unreliable** and are subject to significant variation between years; 3) Rates may not provide statistically valid methods for comparison, particularly among smaller children.