

Table 17B
2005 Pediatric Nutrition Surveillance (1)
Sutter
Anemia Indicators by Race/Ethnicity and Age
Children Aged < 5 Years (2)

Page: 1
Run Date: 11 / 30 / 06

State: California

County: 051

Anemia Indicators by Race/Ethnicity and Age (3, 4)					
		Anemia Low Hb/Hct %			
		Number		Number	Anemia Low Hb/Hct %
White, Not Hispanic					
6 - 11 Months	89		*		
12 - 17 Months	70		*		
18 - 23 Months	29		*		
24 - 35 Months	91		*		
36 - 59 Months	153	9.2			
Total	432	12.3			
Black, Not Hispanic					
6 - 11 Months	2		*		
12 - 17 Months	2		*		
18 - 23 Months	3		*		
24 - 35 Months	2		*		
36 - 59 Months	10		*		
Total	19		*		
Hispanic					
6 - 11 Months	170	10.6			
12 - 17 Months	126	12.7			
18 - 23 Months	50	*			
24 - 35 Months	140	14.3			
36 - 59 Months	219	9.6			
Total	705	11.3			
American Indian/Alaskan Native					
6 - 11 Months	0		*		
12 - 17 Months	2		*		
18 - 23 Months	1		*		
24 - 35 Months	3		*		
36 - 59 Months	2		*		
Total	8		*		
Asian					
6 - 11 Months	22		*		
12 - 17 Months	28		*		
18 - 23 Months	11		*		
24 - 35 Months	20		*		
36 - 59 Months	60		*		
Total	141	18.4			
Pacific Islander					
6 - 11 Months	0		*		
12 - 17 Months	0		*		
18 - 23 Months	0		*		
24 - 35 Months	0		*		
36 - 59 Months	0		*		
Total	0		*		
Filipino					
6 - 11 Months	0		*		
12 - 17 Months	0		*		
18 - 23 Months	0		*		
24 - 35 Months	1		*		
36 - 59 Months	0		*		
Total	1		*		
All Other					
6 - 11 Months	27		*		
12 - 17 Months	15		*		
18 - 23 Months	15		*		
24 - 35 Months	38		*		
36 - 59 Months	94		*		
Total	189	15.3			

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 17B
2005 Pediatric Nutrition Surveillance (1)
Sutter
Anemia Indicators by Race/Ethnicity and Age
Children Aged < 5 Years (2)

State: California

County: 051

Anemia Indicators by Race/Ethnicity and Age (3, 4)			
	Anemia Low Hb/Hct Number	%	
Total All Race/Ethnic Groups	310	12.3	
6 - 11 Months	243	12.8	
12 - 17 Months	109	11.0	
18 - 23 Months	295	16.3	
24 - 35 Months	538	11.3	
36 - 59 Months	1,495	12.7	
Total	1,495	12.7	

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 17B
2005 Pediatric Nutrition Surveillance (1)
Sutter
Anemia Indicators by Race/Ethnicity and Age
Children Aged 5 to <20 Years (2)

State: California

County: 051

Anemia Indicators by Race/Ethnicity and Age (3, 4)					
	Anemia Low Hb/Hct			Anemia Low Hb/Hct	
	Number	%		Number	%
White, Not Hispanic			Asian		
5 - 8 Years	77	*	5 - 8 Years	27	*
9 - 11 Years	41	*	9 - 11 Years	11	*
12 - 14 Years - Girls	16	*	12 - 14 Years - Girls	5	*
12 - 14 Years - Boys	19	*	12 - 14 Years - Boys	8	*
15 - 19 Years - Girls	10	*	15 - 19 Years - Girls	4	*
15 - 19 Years - Boys	13	*	15 - 19 Years - Boys	3	*
Total	176	13.6	Total	58	*
Black, Not Hispanic			Pacific Islander		
5 - 8 Years	11	*	5 - 8 Years	0	*
9 - 11 Years	1	*	9 - 11 Years	0	*
12 - 14 Years - Girls	0	*	12 - 14 Years - Girls	0	*
12 - 14 Years - Boys	1	*	12 - 14 Years - Boys	0	*
15 - 19 Years - Girls	2	*	15 - 19 Years - Girls	0	*
15 - 19 Years - Boys	3	*	15 - 19 Years - Boys	0	*
Total	18	*	Total	0	*
Hispanic			Filipino		
5 - 8 Years	124	12.1	5 - 8 Years	1	*
9 - 11 Years	45	*	9 - 11 Years	2	*
12 - 14 Years - Girls	23	*	12 - 14 Years - Girls	0	*
12 - 14 Years - Boys	18	*	12 - 14 Years - Boys	0	*
15 - 19 Years - Girls	20	*	15 - 19 Years - Girls	0	*
15 - 19 Years - Boys	15	*	15 - 19 Years - Boys	0	*
Total	245	13.9	Total	3	*
American Indian/Alaskan Native			All Other		
5 - 8 Years	6	*	5 - 8 Years	59	*
9 - 11 Years	3	*	9 - 11 Years	26	*
12 - 14 Years - Girls	1	*	12 - 14 Years - Girls	15	*
12 - 14 Years - Boys	0	*	12 - 14 Years - Boys	12	*
15 - 19 Years - Girls	0	*	15 - 19 Years - Girls	7	*
15 - 19 Years - Boys	1	*	15 - 19 Years - Boys	4	*
Total	11	*	Total	123	19.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 17B
2005 Pediatric Nutrition Surveillance ⁽¹⁾
Sutter
Anemia Indicators by Race/Ethnicity and Age
Children Aged 5 to <20 Years ⁽²⁾

State: California

County: 051

Anemia Indicators by Race/Ethnicity and Age ^(3, 4)			
	Anemia Low Hb/Hct Number	%	
Total All Race/Ethnic Groups	305	17.0	
5 - 8 Years	129	17.1	
9 - 11 Years	60	*	
12 - 14 Years - Girls	58	*	
12 - 14 Years - Boys	43	*	
15 - 19 Years - Girls	39	*	
15 - 19 Years - Boys	634	16.9	
Total			

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.