

Table 6B
2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alameda	6,147	7.1 (46)	9.6 (45)	30,928	4.6 (26)	6.5 (31)	13.1 (27)	9,764	15.9 (22)	16.7 (35)	8,455	13.6 (27)
Alpine	38	* *	* *	140	6.4 (57)	8.6 (54)	16.4 (54)	41	* *	* *	70	* *
Amador	106	4.7 (12)	8.5 (33)	671	4.0 (14)	4.5 (9)	14.5 (40)	217	9.2 (1)	17.1 (39)	286	13.3 (23)
Butte	2,474	5.3 (16)	7.5 (22)	8,750	4.8 (32)	5.6 (21)	11.7 (16)	2,692	15.5 (18)	11.4 (3)	4,902	10.6 (9)
Calaveras	245	6.1 (29)	6.9 (10)	1,501	4.5 (23)	7.1 (37)	15.1 (44)	500	17.8 (51)	16.6 (31)	822	15.5 (41)
City Of Be	20	* *	* *	93	* *	* *	* *	25	* *	* *	36	* *
Colusa	416	4.1 (6)	7.5 (22)	1,761	2.6 (1)	6.5 (31)	12.8 (22)	495	14.1 (7)	16.6 (31)	926	14.8 (35)
Contra Cos	3,649	6.9 (45)	9.6 (45)	16,332	4.7 (28)	7.9 (50)	13.2 (33)	4,389	16.0 (25)	17.3 (41)	4,617	14.3 (32)
Del Norte	351	4.6 (11)	12.8 (56)	1,121	5.4 (41)	4.4 (8)	16.1 (51)	406	20.2 (60)	15.5 (16)	679	6.2 (1)
El Dorado	797	6.1 (29)	7.2 (18)	2,810	4.2 (17)	10.9 (61)	9.1 (5)	807	13.4 (4)	13.3 (8)	1,089	11.0 (11)
Fresno	11,308	6.8 (43)	7.4 (19)	57,661	6.0 (53)	4.0 (4)	16.8 (55)	19,768	16.8 (39)	17.2 (40)	33,753	13.5 (25)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Glenn	198	4.0 (4)	7.1 (15)	1,363	5.4 (41)	5.0 (13)	13.1 (27)	447	15.2 (14)	14.8 (12)	889	16.2 (46)
Humboldt	821	5.7 (20)	8.8 (38)	4,186	5.6 (46)	5.0 (13)	12.6 (20)	1,188	17.1 (42)	15.6 (21)	2,104	11.6 (12)
Imperial	2,378	5.6 (18)	8.7 (36)	8,759	3.1 (2)	7.2 (39)	13.1 (27)	3,195	15.7 (20)	16.6 (31)	5,109	8.6 (3)
Inyo	61	* *	* *	504	5.8 (51)	3.4 (1)	13.5 (37)	171	18.1 (52)	14.6 (11)	280	16.8 (48)
Kern	10,024	6.5 (37)	8.0 (29)	43,426	5.1 (37)	6.0 (24)	15.4 (46)	14,672	14.8 (11)	16.3 (30)	21,676	9.6 (6)
Kings	527	4.2 (10)	7.6 (27)	5,071	4.9 (34)	3.9 (3)	19.5 (63)	1,755	17.7 (50)	20.9 (58)	3,116	26.1 (61)
L.A. East	21,001	6.1 (29)	6.9 (10)	50,968	5.7 (50)	4.5 (9)	17.0 (57)	11,232	17.2 (43)	17.4 (43)	18,144	15.4 (39)
L.A. North	7,140	6.3 (35)	7.1 (15)	26,491	6.2 (54)	5.0 (13)	17.0 (57)	5,982	16.1 (29)	16.7 (35)	8,465	12.5 (19)
L.A. Other	20,270	7.1 (46)	7.4 (19)	191,494	4.4 (21)	4.2 (7)	18.3 (61)	82,524	16.4 (31)	18.6 (51)	113,973	13.4 (24)
L.A. South	20,580	6.5 (37)	6.9 (10)	48,240	6.3 (56)	4.1 (5)	18.3 (61)	10,810	17.0 (41)	17.5 (45)	17,705	12.1 (17)
L.A. West	2,889	6.7 (42)	6.4 (6)	8,705	6.8 (61)	4.6 (11)	17.5 (60)	2,221	15.9 (22)	15.8 (25)	2,995	14.2 (31)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Lake	254	9.4 (57)	5.1 (5)	1,118	4.8 (32)	8.1 (51)	11.2 (13)	324	13.6 (5)	13.9 (9)	582	13.2 (22)
Lassen	160	6.3 (35)	5.0 (4)	717	5.0 (36)	9.3 (56)	9.5 (6)	244	13.1 (3)	12.3 (4)	384	12.0 (16)
Long Beach	2,131	6.2 (32)	8.2 (31)	12,158	4.4 (21)	5.5 (20)	17.2 (59)	4,286	18.4 (53)	20.0 (56)	4,972	13.8 (29)
Madera	2,483	4.8 (13)	7.0 (13)	8,125	4.7 (28)	5.7 (23)	14.5 (40)	2,709	16.2 (30)	15.5 (16)	5,408	17.1 (50)
Marin	130	3.8 (1)	11.5 (53)	2,014	3.8 (10)	7.5 (45)	12.9 (25)	600	17.5 (48)	18.7 (52)	1,045	8.8 (4)
Mariposa	27	* *	* *	125	4.0 (14)	6.4 (30)	7.2 (2)	44	* *	* *	70	* *
Mendocino	744	7.1 (46)	9.7 (47)	2,468	3.6 (7)	9.6 (58)	11.6 (15)	659	17.3 (45)	18.1 (48)	891	19.0 (57)
Merced	3,492	5.5 (17)	8.7 (36)	12,277	5.5 (44)	4.6 (11)	15.7 (50)	4,340	16.4 (31)	17.4 (43)	8,254	17.3 (52)
Modoc	51	* *	* *	218	6.4 (57)	7.8 (49)	8.7 (4)	72	* *	* *	122	17.2 (51)
Mono	145	7.6 (52)	3.4 (1)	472	7.0 (62)	11.0 (62)	7.0 (1)	122	16.4 (31)	9.0 (1)	263	17.9 (53)
Monterey	1,051	4.1 (6)	9.9 (49)	6,612	5.5 (44)	3.7 (2)	15.1 (44)	1,774	18.4 (53)	17.5 (45)	3,239	13.1 (21)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Napa	595	5.9 (25)	9.2 (42)	3,578	3.1 (2)	11.5 (63)	9.6 (7)	888	15.2 (14)	18.0 (47)	1,112	24.4 (60)
Nevada	428	8.2 (54)	7.5 (22)	1,300	6.5 (59)	9.8 (59)	10.5 (11)	328	18.6 (56)	15.5 (16)	451	7.3 (2)
Orange	13,973	5.1 (15)	6.8 (7)	30,831	4.1 (16)	6.0 (24)	13.9 (38)	8,581	16.0 (25)	16.1 (27)	27,794	13.5 (25)
Pasadena	349	8.3 (55)	4.6 (3)	2,308	5.6 (46)	5.6 (21)	16.3 (53)	874	16.6 (36)	18.5 (50)	1,341	11.6 (12)
Placer	866	6.6 (41)	11.0 (50)	4,033	5.6 (46)	6.9 (35)	10.4 (9)	1,076	14.6 (9)	14.3 (10)	1,625	15.0 (36)
Plumas	133	6.0 (26)	12.8 (56)	412	7.3 (63)	10.2 (60)	11.4 (14)	100	15.0 (12)	11.0 (2)	159	8.8 (4)
Riverside	14,797	5.8 (22)	7.4 (19)	49,702	5.8 (51)	5.4 (18)	15.5 (47)	13,381	15.0 (12)	15.5 (16)	20,068	15.2 (38)
Sacramento	4,679	6.8 (43)	9.5 (44)	47,469	5.3 (39)	7.2 (39)	13.1 (27)	15,372	16.0 (25)	15.6 (21)	20,916	18.3 (55)
San Benito	123	6.5 (37)	11.4 (51)	1,040	3.7 (8)	7.5 (45)	15.6 (48)	321	19.0 (57)	20.6 (57)	690	12.2 (18)
San Bernar	9,159	6.2 (32)	7.0 (13)	55,941	6.6 (60)	5.4 (18)	16.1 (51)	16,339	15.7 (20)	16.1 (27)	24,193	15.4 (39)
San Diego	17,845	6.2 (32)	8.0 (29)	72,331	4.7 (28)	6.5 (31)	13.3 (34)	22,486	16.4 (31)	15.5 (16)	42,845	12.7 (20)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
San Franci	691	8.1 (53)	4.1 (2)	10,035	4.6 (26)	7.2 (39)	12.1 (17)	3,112	17.5 (48)	17.0 (38)	1,323	11.6 (12)
San Joaqui	8,050	7.5 (51)	8.5 (33)	31,448	5.4 (41)	6.7 (34)	14.5 (40)	10,318	15.9 (22)	16.0 (26)	12,356	14.1 (30)
San Luis O	1,261	4.8 (13)	9.7 (47)	5,892	3.7 (8)	8.4 (53)	10.3 (8)	1,728	14.5 (8)	13.1 (6)	2,234	16.8 (48)
San Mateo	665	5.7 (20)	7.1 (15)	10,306	3.8 (10)	7.3 (44)	12.6 (20)	2,904	16.0 (25)	19.4 (55)	4,002	10.1 (8)
Santa Barb	4,926	12.8 (58)	6.8 (7)	19,514	4.7 (28)	6.0 (24)	13.1 (27)	5,813	17.3 (45)	19.1 (54)	11,086	16.5 (47)
Santa Clar	3,675	5.8 (22)	7.8 (28)	34,065	4.3 (18)	6.0 (24)	13.1 (27)	10,669	15.5 (18)	15.7 (23)	15,832	9.9 (7)
Santa Cruz	672	3.9 (2)	9.4 (43)	2,196	5.3 (39)	7.1 (37)	15.6 (48)	464	19.0 (57)	22.6 (60)	748	11.8 (15)
Shasta	1,663	6.0 (26)	7.5 (22)	5,868	6.2 (54)	5.3 (16)	13.0 (26)	1,854	18.4 (53)	15.3 (15)	3,368	14.3 (32)
Sierra	11	* *	* *	59	* *	* *	* *	19	* *	* *	30	* *
Siskiyou	368	7.1 (46)	6.8 (7)	1,418	5.1 (37)	9.1 (55)	11.1 (12)	452	17.3 (45)	15.0 (13)	812	15.1 (37)
Solano	2,038	8.3 (55)	9.1 (40)	9,570	4.5 (23)	7.7 (48)	12.8 (22)	2,899	16.6 (36)	17.3 (41)	3,195	14.5 (34)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Sonoma	1,453	5.6 (18)	8.8 (38)	9,317	3.3 (4)	7.2 (39)	12.8 (22)	2,599	16.7 (38)	16.1 (27)	2,884	19.6 (58)
Stanislaus	4,559	5.8 (22)	8.6 (35)	19,882	3.8 (10)	7.2 (39)	13.3 (34)	6,494	15.3 (16)	16.6 (31)	13,148	19.8 (59)
Sutter	748	4.1 (6)	11.4 (51)	3,612	3.9 (13)	6.2 (28)	12.3 (18)	1,044	14.0 (6)	15.2 (14)	1,717	16.0 (45)
Tehama	438	3.9 (2)	13.2 (58)	3,176	3.3 (4)	7.6 (47)	10.4 (9)	909	14.6 (9)	13.1 (6)	1,486	18.4 (56)
Trinity	79	* *	* *	350	4.9 (34)	9.4 (57)	13.4 (36)	117	12.0 (2)	21.4 (59)	201	15.9 (43)
Tulare	5,281	6.0 (26)	7.5 (22)	29,884	5.6 (46)	4.1 (5)	16.8 (55)	10,073	16.5 (35)	16.8 (37)	17,818	18.2 (54)
Tuolumne	243	7.4 (50)	9.1 (40)	715	4.3 (18)	6.9 (35)	8.5 (3)	195	19.5 (59)	12.8 (5)	291	13.7 (28)
Ventura	6,570	6.5 (37)	8.2 (31)	24,000	4.5 (23)	5.3 (16)	14.9 (43)	7,344	16.9 (40)	18.1 (48)	14,990	10.7 (10)
Yolo	992	4.0 (4)	12.1 (55)	5,803	3.5 (6)	8.2 (52)	12.3 (18)	1,744	15.3 (16)	15.7 (23)	2,520	15.7 (42)
Yuba	418	4.1 (6)	11.5 (53)	3,063	4.3 (18)	6.2 (28)	14.4 (39)	989	17.2 (43)	18.9 (53)	1,312	15.9 (43)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
California	229,856	6.4	7.7	1,056,397	5.1	5.5	15.4	339,961	16.2	17.0	527,868	14.0
Nation (Prior Year)	1,641,264	9.4	6.6	6,709,486	6.4	4.8	13.9	2,869,075	16.1	14.7	4,381,087	13.6

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alameda	15,657	6.8 (33)	2.6 (36)	17.7 (18)	22.0 (27)	8,263	16.7 (50)
Alpine	62	* *	* *	* *	* *	49	* *
Amador	262	5.0 (9)	3.1 (54)	19.8 (50)	19.8 (10)	207	14.0 (39)
Butte	2,637	8.4 (53)	2.4 (28)	18.1 (29)	20.1 (11)	2,501	10.6 (14)
Calaveras	664	6.2 (23)	1.8 (12)	17.0 (9)	27.0 (58)	614	13.7 (35)
City Of Be	39	* *	* *	* *	* *	33	* *
Colusa	663	4.2 (3)	2.9 (47)	18.3 (32)	27.3 (60)	636	12.7 (28)
Contra Cos	5,848	6.3 (25)	2.0 (16)	18.2 (30)	24.8 (50)	3,247	15.4 (45)
Del Norte	342	7.9 (47)	1.5 (7)	19.9 (52)	22.5 (32)	306	7.2 (3)
El Dorado	836	6.8 (33)	1.8 (12)	16.0 (2)	19.6 (9)	616	6.2 (2)
Fresno	34,936	8.2 (52)	2.4 (28)	17.9 (21)	24.1 (44)	30,842	13.7 (35)
Glenn	345	6.4 (29)	3.5 (58)	16.2 (4)	24.9 (51)	351	12.5 (26)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Humboldt	899	5.7 (16)	2.3 (22)	19.2 (46)	21.7 (22)	655	12.4 (24)
Imperial	2,900	4.3 (4)	2.8 (42)	17.3 (13)	25.5 (53)	2,802	10.6 (14)
Inyo	215	4.7 (6)	2.8 (42)	16.3 (5)	27.4 (61)	139	10.1 (11)
Kern	16,920	5.5 (11)	3.3 (56)	18.0 (26)	23.2 (38)	14,969	11.0 (17)
Kings	1,730	6.6 (31)	1.6 (9)	18.2 (30)	27.1 (59)	1,631	22.3 (60)
L.A. East	21,490	6.4 (29)	2.6 (36)	18.5 (35)	21.5 (21)	21,925	12.6 (27)
L.A. North	15,038	8.8 (56)	2.1 (19)	19.2 (46)	20.3 (13)	15,401	9.0 (10)
L.A. Other	108,672	4.7 (6)	2.3 (22)	18.9 (41)	25.8 (54)	89,785	11.9 (21)
L.A. South	27,696	7.6 (42)	2.2 (21)	19.9 (52)	19.4 (8)	29,469	8.9 (9)
L.A. West	3,917	5.2 (10)	2.5 (33)	17.9 (21)	21.2 (19)	3,496	16.0 (48)
Lake	398	7.3 (39)	2.0 (16)	23.1 (60)	21.1 (18)	323	10.8 (16)
Lassen	409	12.2 (60)	1.2 (5)	16.1 (3)	26.4 (57)	356	13.5 (33)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Long Beach	6,029	4.8 (8)	2.0 (16)	18.9 (41)	24.2 (46)	4,472	14.8 (41)
Madera	3,104	7.0 (36)	2.3 (22)	19.1 (45)	23.0 (35)	3,350	12.8 (30)
Marin	596	12.2 (60)	2.9 (47)	19.8 (50)	23.0 (35)	363	8.0 (6)
Mariposa	65	* *	* *	* *	* *	53	* *
Mendocino	616	7.3 (39)	2.6 (36)	20.1 (54)	20.3 (13)	442	12.9 (31)
Merced	4,489	10.2 (58)	2.1 (19)	18.9 (41)	21.9 (24)	4,568	15.8 (47)
Modoc	106	2.8 (1)	2.8 (42)	17.9 (21)	14.2 (1)	102	4.9 (1)
Mono	108	4.6 (5)	3.7 (59)	17.6 (16)	25.9 (55)	102	12.7 (28)
Monterey	2,576	9.0 (57)	1.3 (6)	20.2 (55)	21.9 (24)	2,608	10.3 (12)
Napa	1,060	7.9 (47)	0.9 (2)	20.2 (55)	23.6 (41)	689	22.5 (61)
Nevada	293	5.5 (11)	3.8 (61)	17.4 (14)	17.7 (4)	197	12.2 (23)
Orange	17,113	7.0 (36)	2.6 (36)	18.6 (36)	20.9 (17)	28,074	13.9 (38)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Pasadena	1,266	5.7 (16)	2.3 (22)	17.8 (20)	24.6 (48)	1,179	7.5 (4)
Placer	1,025	7.8 (45)	2.8 (42)	19.6 (49)	20.1 (11)	645	13.5 (33)
Plumas	145	7.6 (42)	0.7 (1)	17.9 (21)	20.7 (16)	109	8.3 (7)
Riverside	19,214	5.8 (19)	2.9 (47)	17.6 (16)	21.8 (23)	17,605	13.4 (32)
Sacramento	24,131	8.5 (54)	2.6 (36)	17.1 (10)	22.1 (29)	17,512	19.9 (59)
San Benito	322	11.2 (59)	1.6 (9)	24.5 (61)	24.5 (47)	312	15.1 (43)
San Bernar	25,143	6.2 (23)	2.8 (42)	18.0 (26)	22.1 (29)	22,884	13.8 (37)
San Diego	29,446	6.3 (25)	2.5 (33)	18.6 (36)	22.9 (34)	28,962	11.8 (20)
San Franci	6,505	7.9 (47)	3.3 (56)	15.7 (1)	18.3 (6)	2,210	15.2 (44)
San Joaqui	14,941	7.0 (36)	3.0 (51)	17.1 (10)	22.0 (27)	7,884	17.2 (53)
San Luis O	1,560	7.8 (45)	2.4 (28)	20.3 (57)	23.0 (35)	1,042	12.1 (22)
San Mateo	3,362	7.5 (41)	1.6 (9)	21.0 (59)	24.7 (49)	1,841	10.5 (13)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Santa Barb	4,934	7.7 (44)	1.9 (14)	19.5 (48)	25.9 (55)	4,670	12.4 (24)
Santa Clar	15,372	6.8 (33)	3.0 (51)	17.9 (21)	22.5 (32)	8,405	11.0 (17)
Santa Cruz	871	7.9 (47)	0.9 (2)	20.3 (57)	22.3 (31)	812	7.8 (5)
Shasta	1,709	6.3 (25)	1.5 (7)	18.9 (41)	18.3 (6)	1,415	16.6 (49)
Sierra	45	* *	* *	* *	* *	46	* *
Siskiyou	545	4.0 (2)	3.7 (59)	17.2 (12)	18.0 (5)	472	15.5 (46)
Solano	2,929	5.6 (13)	2.4 (28)	18.3 (32)	23.3 (39)	1,491	17.6 (55)
Sonoma	2,463	8.0 (51)	2.5 (33)	18.4 (34)	23.5 (40)	1,213	14.2 (40)
Stanislaus	7,621	5.7 (16)	2.4 (28)	18.0 (26)	24.1 (44)	8,038	17.8 (57)
Sutter	720	5.8 (19)	3.1 (54)	18.6 (36)	20.4 (15)	646	18.6 (58)
Tehama	784	5.6 (13)	1.9 (14)	16.8 (7)	21.3 (20)	689	17.7 (56)
Trinity	167	6.0 (22)	3.0 (51)	17.4 (14)	16.2 (2)	146	17.1 (51)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Tulare	13,106	6.3 (25)	2.6 (36)	18.6 (36)	23.6 (41)	12,890	14.8 (41)
Tuolumne	288	6.6 (31)	1.0 (4)	17.7 (18)	16.3 (3)	256	11.7 (19)
Ventura	6,240	5.8 (19)	2.3 (22)	18.8 (40)	23.9 (43)	6,113	8.6 (8)
Yolo	1,957	5.6 (13)	2.3 (22)	16.8 (7)	24.9 (51)	1,200	17.2 (53)
Yuba	771	8.7 (55)	2.9 (47)	16.5 (6)	21.9 (24)	563	17.1 (51)
California	486,312	6.5	2.5	18.4	23.1	424,886	12.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 - 2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alameda	14,817	7.2 (46)	10.1 (52)	79,415	4.8 (33)	6.5 (33)	13.5 (27)	27,292	16.1 (26)	16.4 (27)	23,888	13.9 (35)
Alpine	101	7.9 (55)	7.9 (24)	437	6.9 (66)	5.9 (26)	16.0 (50)	158	20.3 (64)	20.9 (62)	237	13.5 (30)
Amador	397	7.6 (49)	7.8 (23)	2,101	4.7 (28)	5.5 (20)	14.9 (41)	709	13.8 (6)	15.8 (20)	929	11.9 (15)
Butte	6,024	6.7 (32)	7.7 (20)	22,504	5.0 (39)	6.0 (27)	11.8 (13)	7,977	15.4 (14)	12.6 (5)	13,875	10.4 (7)
Calaveras	555	7.6 (49)	8.5 (34)	3,515	5.6 (49)	6.4 (31)	16.3 (54)	1,378	16.3 (31)	16.6 (30)	2,081	14.6 (40)
City Of Be	53	* *	* *	257	3.9 (10)	7.0 (39)	17.1 (57)	71	* *	* *	126	19.0 (62)
Colusa	988	4.5 (2)	7.5 (16)	4,308	3.3 (2)	7.9 (54)	13.3 (24)	1,429	14.3 (8)	17.8 (42)	2,406	15.6 (48)
Contra Cos	8,317	7.0 (38)	10.1 (52)	41,856	4.9 (38)	7.4 (45)	13.9 (33)	13,064	16.7 (43)	17.3 (39)	12,979	13.0 (25)
Del Norte	732	4.8 (4)	13.0 (62)	2,869	4.8 (33)	4.2 (5)	15.9 (48)	1,114	19.2 (63)	18.0 (48)	1,829	6.8 (1)
El Dorado	1,899	6.4 (23)	7.7 (20)	7,272	4.2 (17)	11.2 (65)	9.6 (4)	2,340	14.2 (7)	13.5 (7)	3,336	10.2 (6)
Fresno	27,621	7.0 (38)	7.6 (19)	147,368	5.7 (51)	4.0 (2)	16.6 (55)	57,304	16.4 (33)	17.1 (35)	94,985	13.0 (25)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 - 2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Glenn	606	5.4 (10)	11.2 (58)	3,482	4.8 (33)	5.2 (17)	14.3 (38)	1,266	15.9 (22)	16.7 (32)	2,374	14.8 (41)
Humboldt	2,051	7.1 (41)	11.2 (58)	10,893	5.2 (41)	5.5 (20)	13.0 (22)	3,616	17.4 (54)	15.5 (16)	5,913	12.1 (16)
Imperial	5,652	6.4 (23)	8.2 (27)	21,809	3.2 (1)	6.9 (37)	14.4 (39)	8,649	16.2 (29)	17.8 (42)	13,794	7.4 (2)
Inyo	225	9.8 (62)	8.0 (26)	1,379	5.5 (47)	4.6 (12)	15.4 (44)	482	16.2 (29)	17.8 (42)	777	14.4 (38)
Kern	24,142	6.9 (36)	8.5 (34)	109,401	4.8 (33)	5.8 (23)	16.0 (50)	41,626	15.6 (17)	17.2 (37)	61,156	10.1 (5)
Kings	1,629	5.0 (6)	9.2 (41)	12,773	4.7 (28)	4.4 (9)	19.9 (66)	4,866	16.8 (44)	21.5 (63)	8,224	25.5 (65)
L.A. East	54,041	6.5 (27)	7.4 (14)	136,731	5.9 (57)	4.3 (8)	17.5 (59)	36,073	16.6 (41)	17.9 (46)	58,074	15.2 (45)
L.A. North	19,695	6.5 (27)	7.1 (8)	76,615	6.2 (62)	4.7 (13)	17.4 (58)	21,172	17.0 (48)	19.4 (58)	30,387	12.9 (23)
L.A. Other	49,330	7.6 (49)	7.5 (16)	515,008	4.2 (17)	4.2 (5)	18.2 (64)	242,945	16.5 (36)	18.5 (52)	320,963	13.3 (29)
L.A. South	53,064	7.1 (41)	7.3 (13)	137,071	6.1 (60)	4.1 (4)	18.2 (64)	36,928	16.8 (44)	17.2 (37)	60,350	12.8 (21)
L.A. West	9,102	6.6 (31)	6.6 (5)	27,927	6.4 (63)	4.4 (9)	17.7 (61)	7,868	16.4 (33)	17.9 (46)	11,507	15.3 (46)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 - 2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Lake	704	9.2 (58)	7.1 (8)	3,105	3.8 (9)	7.7 (51)	10.6 (9)	1,058	15.8 (20)	13.1 (6)	1,864	12.8 (21)
Lassen	423	6.9 (36)	7.1 (8)	1,871	4.5 (23)	9.8 (61)	10.2 (8)	697	12.2 (2)	11.9 (4)	1,022	11.3 (13)
Long Beach	6,480	7.3 (47)	8.3 (30)	34,007	4.7 (28)	5.3 (19)	17.6 (60)	13,231	17.2 (50)	19.8 (60)	15,762	13.5 (30)
Madera	5,522	6.0 (16)	7.5 (16)	19,460	4.6 (25)	4.8 (14)	15.4 (44)	7,285	16.9 (46)	16.1 (23)	14,154	17.3 (56)
Marin	354	4.8 (4)	8.5 (34)	6,695	4.0 (12)	7.8 (53)	11.9 (16)	2,130	16.5 (36)	18.7 (53)	3,654	8.1 (3)
Mariposa	77	* *	* *	453	4.2 (17)	9.3 (58)	8.8 (2)	177	12.4 (4)	10.2 (3)	277	14.4 (38)
Mendocino	2,163	7.7 (53)	9.7 (49)	7,667	3.4 (3)	9.2 (56)	12.1 (17)	2,500	18.3 (60)	18.9 (55)	3,067	17.1 (55)
Merced	8,866	6.8 (35)	8.4 (32)	33,819	5.7 (51)	4.2 (5)	16.1 (52)	12,837	17.7 (58)	18.1 (50)	23,972	15.8 (50)
Modoc	120	3.3 (1)	10.8 (56)	533	4.5 (23)	7.9 (54)	9.9 (6)	209	10.5 (1)	9.1 (2)	316	12.3 (20)
Mono	390	9.7 (61)	4.1 (1)	1,221	5.2 (41)	11.4 (66)	7.3 (1)	337	12.2 (2)	8.9 (1)	670	15.1 (44)
Monterey	3,300	5.0 (6)	10.1 (52)	21,479	5.6 (49)	3.6 (1)	15.6 (47)	6,104	18.9 (61)	18.0 (48)	10,854	13.0 (25)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 - 2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Napa	1,559	6.4 (23)	9.3 (42)	8,914	3.4 (3)	9.8 (61)	10.8 (10)	2,707	16.1 (26)	16.0 (22)	3,501	22.4 (64)
Nevada	1,046	7.1 (41)	6.7 (6)	3,602	5.2 (41)	9.8 (61)	9.3 (3)	1,003	17.2 (50)	13.9 (9)	1,499	11.1 (11)
Orange	36,041	5.8 (14)	7.4 (14)	87,327	6.0 (59)	5.6 (22)	16.9 (56)	29,623	15.8 (20)	19.0 (56)	90,212	13.6 (32)
Pasadena	1,377	9.4 (59)	4.3 (2)	7,143	4.7 (28)	6.6 (34)	14.0 (34)	2,836	14.8 (9)	14.9 (13)	4,562	10.9 (10)
Placer	2,233	6.3 (20)	9.6 (48)	9,924	5.7 (51)	7.5 (48)	11.3 (12)	2,993	14.9 (10)	14.9 (13)	4,816	13.9 (35)
Plumas	302	9.6 (60)	8.3 (30)	1,161	6.1 (60)	10.4 (64)	12.6 (21)	378	13.2 (5)	16.4 (27)	508	10.4 (7)
Riverside	35,567	6.4 (23)	7.7 (20)	126,100	5.7 (51)	5.1 (16)	15.9 (48)	39,042	15.1 (12)	15.7 (18)	57,558	15.8 (50)
Sacramento	11,970	7.0 (38)	9.5 (47)	121,092	5.2 (41)	6.9 (37)	13.6 (30)	44,444	16.0 (25)	15.7 (18)	59,753	16.5 (54)
San Benito	462	5.0 (6)	13.0 (62)	3,587	3.7 (6)	5.8 (23)	17.9 (63)	1,262	19.1 (62)	23.6 (64)	2,633	15.9 (52)
San Bernar	26,972	7.1 (41)	7.2 (12)	146,531	6.6 (64)	5.2 (17)	16.1 (52)	48,691	15.9 (22)	16.3 (26)	70,778	16.1 (53)
San Diego	42,254	6.5 (27)	8.8 (39)	177,089	4.6 (25)	6.4 (31)	13.3 (24)	60,819	16.1 (26)	15.5 (16)	108,690	12.2 (18)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 - 2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
San Franci	1,975	7.4 (48)	5.8 (3)	24,394	4.3 (20)	7.1 (40)	12.2 (19)	8,344	17.1 (49)	17.0 (33)	3,819	12.2 (18)
San Joaqui	19,051	8.2 (56)	8.2 (27)	76,269	5.4 (46)	6.6 (34)	14.9 (41)	27,774	15.9 (22)	16.4 (27)	30,998	13.0 (25)
San Luis O	3,380	6.7 (32)	9.7 (49)	14,908	3.9 (10)	9.3 (58)	10.8 (10)	4,774	15.5 (15)	13.9 (9)	6,245	15.7 (49)
San Mateo	2,479	7.1 (41)	8.6 (37)	30,588	4.0 (12)	7.2 (41)	13.5 (27)	9,894	17.5 (56)	19.6 (59)	13,449	9.3 (4)
Santa Barb	12,503	14.1 (63)	7.1 (8)	48,742	4.8 (33)	5.8 (23)	14.0 (34)	16,602	17.2 (50)	19.3 (57)	30,496	14.9 (42)
Santa Clar	9,122	6.0 (16)	8.4 (32)	89,070	4.1 (16)	6.1 (28)	13.4 (26)	31,673	15.3 (13)	15.9 (21)	42,463	11.3 (13)
Santa Cruz	2,149	5.5 (11)	9.4 (43)	8,135	5.2 (41)	7.3 (44)	14.0 (34)	2,020	16.6 (41)	20.4 (61)	3,007	14.0 (37)
Shasta	3,978	7.6 (49)	8.8 (39)	14,809	6.6 (64)	4.5 (11)	13.7 (31)	5,346	17.5 (56)	15.1 (15)	9,061	12.1 (16)
Sierra	49	* *	* *	187	5.9 (57)	9.6 (60)	9.6 (4)	55	* *	* *	97	* *
Siskiyou	892	6.5 (27)	6.1 (4)	3,690	5.0 (39)	9.2 (56)	11.8 (13)	1,334	17.2 (50)	16.1 (23)	2,373	17.6 (57)
Solano	5,085	7.8 (54)	9.4 (43)	23,110	4.7 (28)	7.7 (51)	13.1 (23)	7,855	16.5 (36)	17.6 (41)	9,024	11.2 (12)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 - 2006 Pediatric Nutrition Surveillance ⁽¹⁾
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by County ^(3,4)												
County	Birthweight ⁽⁵⁾			Height and Weight						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature ⁽⁶⁾	Under- weight ⁽⁷⁾	Over- weight ⁽⁷⁾		≥ 2 Yrs Over- weight ⁽⁷⁾				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Sonoma	3,990	6.0 (16)	9.4 (43)	24,500	3.4 (3)	7.5 (48)	12.4 (20)	7,892	16.5 (36)	16.1 (23)	8,945	18.7 (60)
Stanislaus	11,739	8.3 (57)	8.2 (27)	52,133	4.0 (12)	7.2 (41)	14.2 (37)	19,301	15.7 (18)	17.1 (35)	38,398	18.3 (58)
Sutter	1,914	6.1 (19)	11.0 (57)	8,765	4.0 (12)	7.4 (45)	12.1 (17)	2,893	15.5 (15)	14.0 (11)	4,638	15.0 (43)
Tehama	1,258	5.5 (11)	12.5 (61)	7,492	3.7 (6)	7.2 (41)	11.8 (13)	2,578	14.9 (10)	14.2 (12)	3,902	19.1 (63)
Trinity	177	5.1 (9)	6.8 (7)	796	5.7 (51)	7.4 (45)	14.6 (40)	283	16.3 (31)	17.0 (33)	473	18.4 (59)
Tulare	13,587	6.7 (32)	7.9 (24)	77,406	5.8 (56)	4.0 (2)	17.7 (61)	29,311	16.9 (46)	17.8 (42)	49,924	18.8 (61)
Tuolumne	522	5.9 (15)	10.2 (55)	2,052	5.5 (47)	6.6 (34)	10.1 (7)	686	17.9 (59)	13.7 (8)	1,077	13.7 (33)
Ventura	15,885	6.3 (20)	8.6 (37)	61,444	4.4 (22)	5.0 (15)	15.4 (44)	21,346	16.4 (33)	18.3 (51)	40,944	10.8 (9)
Yolo	2,350	4.5 (2)	12.2 (60)	14,706	3.7 (6)	7.6 (50)	13.5 (27)	5,097	15.7 (18)	17.5 (40)	7,469	13.8 (34)
Yuba	1,218	6.3 (20)	9.9 (51)	7,300	4.6 (25)	6.3 (30)	13.8 (32)	2,542	17.4 (54)	16.6 (30)	3,540	15.4 (47)
Unknown Counties	1,992	5.6 (13)	9.4 (43)	8,257	4.3 (20)	6.1 (28)	15.1 (43)	3,134	16.5 (36)	18.7 (53)	4,012	12.9 (23)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 - 2006 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
California	584,518	7.0	8.0	2,786,524	5.0	5.4	15.8	1,009,424	16.3	17.3	1,524,666	13.8

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.