

**Table 2C**  
**2007 Pediatric Nutrition Surveillance** (1)  
**California**  
**Summary of Health Indicators**  
**Children Aged < 5 Years** (2)

<b>Summary of Health Indicators</b> (3)						
			California Prevalence		National Prevalence (Prior Year)	
			Number	%	Number	%
<b>Birthweight (4)</b>						
Low	< 2500 g		220,928	6.4	1,842,970	9.2
High	> 4000 g		220,928	7.5	1,842,970	6.6
<b>Height and Weight</b>						
<b>Short Stature (5)</b>						
	< 5th		1,016,899	5.1	7,192,043	6.4
	< 10th		1,016,899	9.0	7,192,043	11.4
<b>Underweight (6)</b>						
	< 5th		1,016,899	5.4	7,192,043	4.7
	< 10th		1,016,899	9.0	7,192,043	8.0
<b>Overweight (6)</b>						
	≥ 95th		1,016,899	15.5	7,192,043	13.8
≥ 2 Yrs Overweight (6)	85th - < 95th		312,190	16.2	2,945,212	16.4
	≥ 95th		312,190	17.4	2,945,212	14.8
<b>Anemia (7)</b>						
Low Hb			443,462	14.8	4,092,770	14.3
Low Hct			58,456	8.9	652,304	10.9
Low Hb/Hct			474,562	14.2	4,521,591	14.0
<b>Breastfeeding (8)</b>						
Ever Breastfed			0	*	1,577,832	60.1
Breastfed At Least 6 Months			0	*	790,341	25.2
Breastfed At Least 12 Months			0	*	875,205	18.1
<b>Smoking in Household</b>						
≤ 2 Hours TV Viewing/ Day (9)			1,146,691	4.1	2,855,916	10.6
			0	*	584,732	73.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

(8) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(9) Children aged 2 years and older included in the analysis.

\* Percentages are not calculated if < 100 records are available for analysis after exclusions.

**Table 2C**  
**2007 Pediatric Nutrition Surveillance** <sup>(1)</sup>  
**California**  
**Summary of Health Indicators**  
**Children Aged 5 to <20 Years** <sup>(2)</sup>

<b>Summary of Health Indicators</b> <sup>(3)</sup>					
		California Prevalence		National Prevalence (Prior Year)	
		Number	%	Number	%
Height and Weight					
Short Stature <sup>(4)</sup>	< 5th	473,184	6.6	0	*
Underweight <sup>(5)</sup>	< 5th	473,184	2.4	0	*
Overweight <sup>(5)</sup>	85th - < 95th	473,184	18.4	0	*
	≥ 95th	473,184	23.1	0	*
Anemia <sup>(6)</sup>					
Low Hb		374,602	13.1	0	*
Low Hct		48,685	11.8	0	*
Low Hb/Hct		400,067	12.9	0	*
Smoking in Household		547,326	5.2	0	*
≤2 Hours TV Viewing/Day <sup>(7)</sup>		0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for height-for-age.

(5) Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - <95th percentile category identifies children at risk of overweight.

(6) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

(7) Children aged 2 years and older included in the analysis.

\* Percentages are not calculated if < 100 records are available for analysis after exclusions.