

Table 6B
2007 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alameda	6,390	6.7 (40)	8.4 (32)	32,350	4.4 (22)	6.6 (31)	13.2 (26)	10,371	16.2 (29)	16.4 (23)	8,652	15.2 (42)
Alpine	30	* *	* *	141	7.8 (62)	7.8 (47)	14.9 (45)	43	* *	* *	74	* *
Amador	143	9.8 (56)	1.4 (1)	813	7.0 (61)	5.4 (18)	13.7 (33)	241	15.8 (23)	13.3 (6)	268	13.4 (30)
Butte	2,217	6.2 (29)	9.7 (42)	8,532	4.3 (20)	6.1 (26)	11.3 (12)	2,680	12.9 (5)	12.5 (2)	5,004	11.4 (12)
Calaveras	159	1.9 (1)	6.9 (15)	1,536	4.6 (26)	5.2 (15)	16.3 (55)	520	15.4 (17)	20.0 (54)	875	14.2 (35)
City Of Be	17	* *	* *	96	* *	* *	* *	31	* *	* *	48	* *
Colusa	367	4.4 (8)	10.9 (51)	1,701	2.4 (1)	7.6 (44)	14.3 (40)	471	14.9 (11)	17.2 (37)	838	18.9 (53)
Contra Cos	3,398	6.6 (38)	9.5 (41)	16,650	4.6 (26)	7.6 (44)	12.7 (20)	4,354	16.5 (32)	17.1 (34)	4,317	14.6 (37)
Del Norte	289	3.1 (2)	13.1 (55)	1,088	5.3 (42)	5.0 (13)	14.8 (43)	353	19.0 (56)	17.6 (40)	589	8.5 (2)
El Dorado	678	5.6 (17)	11.8 (53)	2,936	3.9 (12)	10.9 (60)	9.5 (4)	846	12.4 (4)	13.0 (4)	1,166	11.3 (11)
Fresno	10,812	7.0 (43)	7.7 (26)	59,930	6.0 (54)	4.1 (4)	17.1 (57)	20,198	16.8 (36)	18.0 (45)	35,893	15.1 (40)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2007 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Glenn	204	3.9 (7)	11.3 (52)	1,420	3.6 (7)	5.4 (18)	13.9 (36)	457	16.8 (36)	15.8 (17)	979	19.3 (54)
Humboldt	706	6.2 (29)	9.9 (43)	4,288	5.0 (37)	5.9 (24)	12.4 (18)	1,232	15.5 (18)	14.2 (7)	2,169	12.4 (23)
Imperial	2,335	5.7 (19)	7.9 (27)	8,510	2.8 (2)	7.2 (35)	13.7 (33)	3,004	15.3 (15)	17.7 (42)	4,898	5.3 (1)
Inyo	25	* *	* *	436	5.3 (42)	4.1 (4)	15.8 (52)	156	20.5 (59)	16.0 (20)	271	11.8 (18)
Kern	10,308	7.1 (45)	7.9 (27)	44,326	5.0 (37)	5.3 (16)	15.4 (47)	14,429	14.9 (11)	15.9 (18)	21,315	10.7 (6)
Kings	570	5.8 (23)	10.5 (49)	4,536	5.5 (45)	3.4 (2)	21.4 (63)	1,534	18.9 (55)	21.3 (57)	2,947	26.0 (60)
L.A. East	20,512	5.8 (23)	6.6 (9)	48,640	6.4 (57)	4.0 (3)	17.9 (58)	10,179	16.9 (39)	17.5 (39)	16,290	15.7 (44)
L.A. North	7,210	6.1 (26)	5.7 (6)	25,833	6.0 (54)	4.8 (12)	16.1 (54)	5,317	15.9 (24)	17.0 (32)	7,268	11.6 (15)
L.A. Other	20,729	7.2 (47)	6.9 (15)	178,930	4.6 (26)	4.1 (4)	18.6 (60)	71,872	16.4 (31)	19.1 (52)	104,321	13.2 (27)
L.A. South	19,442	6.2 (29)	6.4 (8)	46,664	6.6 (58)	4.1 (4)	18.7 (61)	10,077	17.2 (47)	18.9 (49)	16,044	12.0 (20)
L.A. West	2,097	6.1 (26)	7.1 (20)	6,787	7.8 (62)	4.2 (8)	19.6 (62)	1,590	17.7 (52)	17.1 (34)	2,238	13.2 (27)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2007 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Lake	169	6.5 (36)	10.1 (46)	1,076	4.1 (19)	9.5 (57)	13.2 (26)	346	13.9 (7)	15.6 (13)	575	9.2 (5)
Lassen	167	5.4 (15)	5.4 (4)	806	4.7 (30)	10.3 (59)	9.7 (5)	257	11.3 (1)	14.4 (8)	362	13.0 (26)
Long Beach	2,301	7.4 (51)	6.8 (13)	10,884	5.5 (45)	4.2 (8)	18.1 (59)	3,666	17.9 (53)	20.9 (56)	4,327	13.7 (33)
Madera	2,625	5.1 (11)	7.4 (22)	9,574	5.2 (41)	5.3 (16)	14.5 (41)	3,178	15.7 (21)	15.9 (18)	6,413	20.0 (57)
Marin	122	5.7 (19)	13.9 (56)	2,308	3.6 (7)	7.2 (35)	13.4 (28)	679	16.9 (39)	18.3 (47)	1,024	10.8 (7)
Mariposa	22	* *	* *	124	5.6 (49)	8.1 (49)	14.5 (41)	40	* *	* *	61	* *
Mendocino	765	6.3 (34)	6.9 (15)	2,561	3.6 (7)	8.6 (53)	13.0 (23)	760	20.1 (58)	21.8 (58)	943	16.8 (48)
Merced	3,347	5.3 (14)	9.3 (39)	12,396	5.5 (45)	4.3 (10)	15.6 (49)	4,000	17.0 (42)	17.6 (40)	7,450	18.2 (51)
Modoc	46	* *	* *	255	3.5 (6)	12.2 (62)	9.0 (3)	83	* *	* *	133	9.0 (3)
Mono	114	6.1 (26)	5.3 (3)	436	5.5 (45)	12.4 (63)	8.0 (1)	114	13.2 (6)	7.9 (1)	268	21.3 (59)
Monterey	1,412	3.7 (6)	9.4 (40)	7,136	5.9 (53)	3.3 (1)	15.4 (47)	1,546	17.1 (43)	18.2 (46)	3,294	12.5 (25)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2007 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Napa	414	7.2 (47)	9.9 (43)	3,502	3.9 (12)	9.6 (58)	10.3 (9)	834	16.1 (26)	16.8 (30)	1,204	26.7 (61)
Nevada	335	5.7 (19)	10.1 (46)	1,125	4.0 (16)	10.9 (60)	8.2 (2)	250	15.6 (20)	16.4 (23)	440	11.1 (9)
Orange	7,268	5.1 (11)	6.9 (15)	26,152	4.4 (22)	6.1 (26)	13.5 (29)	7,697	15.9 (24)	16.4 (23)	11,152	13.6 (32)
Pasadena	305	6.2 (29)	2.6 (2)	1,925	6.8 (59)	7.2 (35)	14.2 (38)	612	17.2 (47)	18.6 (48)	1,088	11.6 (15)
Placer	975	7.0 (43)	6.8 (13)	4,267	5.6 (49)	7.2 (35)	12.3 (16)	1,076	14.1 (8)	16.0 (20)	1,801	17.2 (50)
Plumas	111	5.4 (15)	5.4 (4)	395	5.8 (51)	9.4 (56)	9.9 (6)	122	12.3 (3)	13.1 (5)	166	9.0 (3)
Riverside	14,806	6.3 (34)	7.0 (19)	42,148	6.0 (54)	5.4 (18)	14.8 (43)	10,083	14.2 (9)	15.3 (11)	15,181	14.9 (38)
Sacramento	4,920	6.2 (29)	9.1 (38)	46,387	4.8 (34)	7.8 (47)	12.5 (19)	14,538	15.3 (15)	15.4 (12)	19,580	18.6 (52)
San Benito	111	7.2 (47)	16.2 (57)	866	4.8 (34)	7.3 (41)	15.7 (50)	240	15.0 (13)	22.1 (59)	527	12.0 (20)
San Bernar	10,060	6.7 (40)	6.7 (11)	46,845	6.9 (60)	5.4 (18)	16.0 (53)	12,088	15.7 (21)	16.6 (28)	17,573	15.5 (43)
San Diego	17,633	6.6 (38)	8.5 (33)	68,061	4.8 (34)	6.1 (26)	13.9 (36)	19,603	16.5 (32)	16.4 (23)	37,537	11.8 (18)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2007 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
San Franci	1,074	7.4 (51)	6.6 (9)	10,886	4.7 (30)	6.7 (33)	12.3 (16)	3,363	17.5 (51)	16.6 (28)	1,606	11.7 (17)
San Joaqui	7,650	7.2 (47)	7.4 (22)	31,898	5.1 (39)	7.0 (34)	14.2 (38)	10,216	14.7 (10)	16.0 (20)	10,777	14.9 (38)
San Luis O	950	7.1 (45)	8.5 (33)	5,948	3.9 (12)	8.3 (51)	10.5 (10)	1,682	16.8 (36)	15.2 (10)	2,463	16.0 (46)
San Mateo	698	6.9 (42)	5.7 (6)	10,735	4.0 (16)	8.3 (51)	12.0 (14)	3,001	17.1 (43)	17.9 (44)	4,431	10.8 (7)
Santa Barb	4,336	9.9 (57)	8.1 (30)	19,902	4.7 (30)	6.6 (31)	13.6 (31)	6,030	16.9 (39)	19.1 (52)	11,428	16.4 (47)
Santa Clar	3,871	5.0 (10)	7.6 (25)	33,207	4.5 (24)	5.9 (24)	13.5 (29)	9,736	16.2 (29)	17.2 (37)	13,205	11.1 (9)
Santa Cruz	832	3.6 (5)	10.2 (48)	2,770	5.1 (39)	6.5 (29)	15.2 (46)	458	19.9 (57)	20.5 (55)	865	13.4 (30)
Shasta	1,791	8.8 (55)	9.0 (37)	5,946	5.8 (51)	5.4 (18)	12.9 (22)	1,820	17.4 (49)	15.7 (15)	3,223	12.4 (23)
Sierra	10	* *	* *	40	* *	* *	* *	13	* *	* *	17	* *
Siskiyou	364	7.7 (54)	7.4 (22)	1,495	4.7 (30)	9.3 (55)	9.9 (6)	450	18.4 (54)	14.7 (9)	815	15.7 (44)
Solano	2,016	7.4 (51)	7.9 (27)	9,486	4.5 (24)	7.3 (41)	12.8 (21)	3,004	17.1 (43)	17.7 (42)	3,417	14.0 (34)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2007 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Sonoma	1,970	5.2 (13)	8.9 (35)	9,787	3.7 (11)	7.3 (41)	13.0 (23)	2,692	16.7 (35)	17.1 (34)	2,796	17.1 (49)
Stanislaus	4,288	5.8 (23)	7.3 (21)	18,940	4.0 (16)	7.2 (35)	13.7 (33)	5,879	16.1 (26)	16.5 (27)	11,574	21.2 (58)
Sutter	769	3.5 (4)	10.5 (49)	3,871	3.4 (4)	6.5 (29)	12.2 (15)	1,122	15.5 (18)	15.7 (15)	1,925	11.4 (12)
Tehama	432	3.2 (3)	8.3 (31)	3,575	3.0 (3)	8.7 (54)	10.1 (8)	1,009	12.2 (2)	12.7 (3)	1,495	13.3 (29)
Trinity	43	* *	* *	234	3.4 (4)	8.1 (49)	10.7 (11)	68	* *	* *	123	12.2 (22)
Tulare	5,380	5.6 (17)	6.7 (11)	29,798	5.4 (44)	4.5 (11)	16.3 (55)	9,955	16.6 (34)	17.0 (32)	17,471	19.7 (55)
Tuolumne	84	* *	* *	433	3.9 (12)	7.2 (35)	11.3 (12)	109	17.4 (49)	15.6 (13)	161	19.9 (56)
Ventura	6,372	5.7 (19)	8.9 (35)	23,571	4.6 (26)	5.0 (13)	15.7 (50)	7,000	17.1 (43)	18.9 (49)	14,840	11.5 (14)
Yolo	806	4.7 (9)	10.0 (45)	5,562	3.6 (7)	7.6 (44)	13.6 (31)	1,730	16.1 (26)	19.0 (51)	2,602	15.1 (40)
Yuba	526	6.5 (36)	12.0 (54)	3,442	4.3 (20)	5.7 (23)	13.1 (25)	1,086	15.2 (14)	16.8 (30)	1,765	14.3 (36)
Unknown Counties	0	* *	* *	1	* *	* *	* *	0	* *	* *	0	* *

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2007 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
California	220,928	6.4	7.5	1,016,899	5.1	5.4	15.5	312,190	16.2	17.4	474,562	14.2
Nation (Prior Year)	1,842,970	9.2	6.6	7,192,043	6.4	4.7	13.8	2,945,212	16.4	14.8	4,521,591	14.0

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2007 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alameda	17,488	6.9 (34)	2.7 (45)	17.5 (12)	21.8 (21)	9,522	18.2 (53)
Alpine	61	* *	* *	* *	* *	51	* *
Amador	278	6.8 (32)	1.8 (12)	14.7 (3)	19.4 (5)	185	15.1 (37)
Butte	2,669	7.9 (45)	1.7 (9)	18.9 (39)	21.0 (13)	2,514	12.0 (18)
Calaveras	726	6.7 (29)	2.6 (41)	22.3 (58)	24.4 (44)	656	15.5 (39)
City Of Be	36	* *	* *	* *	* *	35	* *
Colusa	574	4.0 (2)	1.6 (7)	18.3 (29)	23.9 (41)	518	14.9 (36)
Contra Cos	6,016	7.9 (45)	2.0 (23)	18.3 (29)	25.4 (50)	2,537	17.4 (48)
Del Norte	274	7.7 (43)	1.8 (12)	21.2 (56)	21.9 (22)	229	9.6 (8)
El Dorado	906	6.1 (16)	3.1 (54)	14.2 (2)	16.8 (1)	608	10.7 (12)
Fresno	35,973	7.9 (45)	2.3 (36)	18.2 (27)	24.4 (44)	32,536	14.8 (35)
Glenn	331	7.9 (45)	1.8 (12)	19.3 (47)	25.7 (54)	342	12.6 (25)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2007 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Humboldt	946	6.6 (27)	2.0 (23)	17.7 (16)	23.6 (38)	723	15.5 (39)
Imperial	2,818	4.5 (3)	3.4 (58)	17.7 (16)	24.8 (46)	2,887	8.8 (4)
Inyo	232	9.1 (56)	1.7 (9)	18.1 (25)	21.6 (18)	146	13.0 (26)
Kern	18,138	5.2 (7)	3.1 (54)	17.5 (12)	24.0 (43)	15,648	12.3 (21)
Kings	1,487	5.6 (11)	3.1 (54)	17.5 (12)	25.2 (48)	1,468	22.0 (57)
L.A. East	21,481	6.3 (20)	2.6 (41)	18.0 (23)	21.2 (15)	26,953	10.6 (11)
L.A. North	14,718	10.3 (59)	2.2 (30)	19.2 (45)	20.4 (8)	14,345	8.1 (3)
L.A. Other	102,738	4.9 (6)	2.3 (36)	18.7 (34)	25.8 (57)	84,580	11.9 (17)
L.A. South	29,713	9.0 (55)	2.1 (27)	19.3 (47)	18.3 (3)	30,205	9.1 (5)
L.A. West	3,046	6.0 (14)	2.9 (50)	19.0 (42)	20.8 (10)	2,688	13.4 (27)
Lake	383	7.3 (38)	1.3 (2)	15.4 (5)	21.4 (17)	306	14.4 (32)
Lassen	416	13.0 (60)	1.9 (19)	18.8 (38)	26.4 (59)	352	10.5 (10)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2007 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Long Beach	5,728	6.1 (16)	1.8 (12)	19.1 (43)	25.4 (50)	4,328	16.1 (42)
Madera	4,004	7.4 (40)	2.6 (41)	18.7 (34)	23.4 (37)	4,207	13.4 (27)
Marin	691	6.9 (34)	1.6 (7)	17.9 (21)	25.3 (49)	359	11.4 (15)
Mariposa	63	* *	* *	* *	* *	53	* *
Mendocino	703	4.7 (5)	0.7 (1)	24.2 (60)	23.9 (41)	483	11.6 (16)
Merced	4,444	9.2 (57)	2.0 (23)	18.7 (34)	23.0 (32)	4,099	17.3 (47)
Modoc	129	2.3 (1)	3.1 (54)	17.8 (20)	20.9 (12)	127	7.1 (1)
Mono	114	7.0 (36)	1.8 (12)	12.3 (1)	17.5 (2)	103	25.2 (58)
Monterey	2,238	7.5 (41)	1.3 (2)	20.0 (51)	22.7 (28)	2,298	9.5 (7)
Napa	1,058	6.3 (20)	1.5 (5)	19.3 (47)	26.0 (58)	721	19.0 (55)
Nevada	260	7.7 (43)	1.9 (19)	22.3 (58)	20.8 (10)	180	14.4 (32)
Orange	16,354	6.3 (20)	2.4 (40)	18.5 (32)	21.1 (14)	15,662	14.1 (31)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2007 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Pasadena	1,004	5.3 (9)	2.3 (36)	17.6 (15)	22.9 (31)	1,000	9.4 (6)
Placer	972	7.9 (45)	3.4 (58)	17.7 (16)	20.2 (7)	652	18.1 (51)
Plumas	134	6.0 (14)	2.2 (30)	14.9 (4)	23.1 (34)	89	* *
Riverside	14,655	5.7 (13)	2.8 (47)	18.1 (25)	22.3 (25)	13,600	12.4 (23)
Sacramento	22,255	7.9 (45)	2.7 (45)	17.4 (10)	21.2 (15)	14,799	19.4 (56)
San Benito	291	10.0 (58)	1.4 (4)	20.6 (53)	26.5 (60)	293	16.7 (45)
San Bernar	20,179	6.4 (24)	2.8 (47)	17.9 (21)	22.0 (23)	18,960	13.4 (27)
San Diego	27,221	6.7 (29)	2.1 (27)	18.9 (39)	22.8 (29)	26,056	10.8 (13)
San Franci	6,905	7.6 (42)	2.8 (47)	16.8 (8)	18.7 (4)	2,425	16.6 (44)
San Joaqui	15,930	6.6 (27)	3.0 (51)	17.2 (9)	22.6 (27)	7,634	15.5 (39)
San Luis O	1,558	6.1 (16)	2.2 (30)	18.9 (39)	21.6 (18)	1,264	12.0 (18)
San Mateo	3,515	7.9 (45)	1.5 (5)	20.8 (54)	25.7 (54)	1,780	13.5 (30)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2007 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Santa Barb	5,260	8.0 (53)	1.7 (9)	20.2 (52)	24.9 (47)	4,992	12.1 (20)
Santa Clar	15,135	7.0 (36)	3.0 (51)	17.7 (16)	22.8 (29)	8,014	12.5 (24)
Santa Cruz	760	8.6 (54)	2.2 (30)	19.6 (50)	23.0 (32)	687	7.4 (2)
Shasta	1,591	7.9 (45)	1.8 (12)	18.2 (27)	19.7 (6)	1,332	14.4 (32)
Sierra	31	* *	* *	* *	* *	23	* *
Siskiyou	546	5.3 (9)	2.0 (23)	17.4 (10)	23.3 (36)	470	11.3 (14)
Solano	3,216	6.7 (29)	2.2 (30)	19.2 (45)	25.4 (50)	1,690	18.0 (49)
Sonoma	2,678	6.1 (16)	1.9 (19)	20.8 (54)	22.1 (24)	1,173	15.3 (38)
Stanislaus	7,618	5.2 (7)	2.2 (30)	18.0 (23)	23.8 (40)	7,487	18.0 (49)
Sutter	872	5.6 (11)	3.6 (60)	16.6 (7)	22.4 (26)	756	18.1 (51)
Tehama	829	4.5 (3)	1.8 (12)	18.7 (34)	20.6 (9)	616	16.1 (42)
Trinity	97	* *	* *	* *	* *	77	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2007 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Tulare	13,264	6.8 (32)	2.3 (36)	18.3 (29)	25.5 (53)	13,067	17.2 (46)
Tuolumne	108	6.5 (26)	1.9 (19)	21.3 (57)	23.1 (34)	86	* *
Ventura	6,428	6.3 (20)	2.1 (27)	18.6 (33)	23.6 (38)	6,477	10.0 (9)
Yolo	2,003	6.4 (24)	2.6 (41)	15.8 (6)	25.7 (54)	1,166	18.6 (54)
Yuba	894	7.3 (38)	3.0 (51)	19.1 (43)	21.7 (20)	747	12.3 (21)
Unknown Counties	1	* *	* *	* *	* *	1	* *
California	473,184	6.6	2.4	18.4	23.1	400,067	12.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.