

Table 6B
2006 - 2008 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alameda	18,216	6.7 (45)	9.0 (42)	94,351	4.5 (22)	6.6 (33)	13.0 (28)	29,996	16.2 (30)	16.6 (33)	25,451	14.0 (32)
Alpine	88	* *	* *	372	6.7 (62)	8.6 (54)	14.0 (39)	112	22.3 (63)	18.8 (54)	188	16.5 (49)
Amador	329	7.0 (47)	4.6 (2)	2,241	5.4 (44)	5.6 (22)	14.1 (40)	675	15.1 (16)	14.2 (10)	773	13.5 (29)
Butte	7,077	5.7 (19)	8.5 (37)	26,165	4.5 (22)	5.9 (23)	11.5 (14)	8,317	15.0 (15)	12.3 (3)	15,244	11.5 (10)
Calaveras	609	4.8 (7)	8.2 (32)	4,572	4.3 (16)	6.1 (28)	15.9 (53)	1,562	16.4 (36)	18.8 (54)	2,561	13.9 (30)
City Of Be	69	* *	* *	281	4.3 (16)	6.0 (25)	11.7 (16)	77	* *	* *	123	24.4 (62)
Colusa	1,120	4.8 (7)	9.8 (51)	4,960	2.9 (1)	7.1 (38)	14.1 (40)	1,385	14.0 (9)	17.3 (42)	2,464	18.2 (53)
Contra Cos	10,841	6.6 (42)	9.8 (51)	49,257	4.4 (20)	8.0 (50)	12.7 (22)	12,874	16.3 (32)	16.8 (36)	12,217	14.9 (39)
Del Norte	765	4.4 (2)	13.3 (61)	3,007	5.5 (48)	4.5 (8)	15.4 (49)	1,066	19.5 (62)	16.3 (27)	1,807	7.1 (2)
El Dorado	2,218	5.7 (19)	8.3 (34)	8,752	4.0 (12)	11.1 (63)	9.3 (3)	2,549	13.1 (4)	13.1 (5)	3,624	12.0 (12)
Fresno	33,570	7.1 (49)	7.2 (14)	178,875	5.9 (56)	4.1 (3)	16.9 (59)	60,982	16.9 (47)	17.5 (45)	105,810	14.4 (35)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 - 2008 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Glenn	750	4.4 (2)	10.5 (54)	4,492	4.9 (36)	4.5 (8)	15.1 (46)	1,458	15.8 (23)	16.9 (37)	3,018	18.3 (55)
Humboldt	2,166	6.0 (27)	9.4 (47)	12,788	5.1 (38)	5.4 (17)	12.8 (24)	3,636	16.5 (37)	15.6 (21)	6,402	11.8 (11)
Imperial	7,474	6.3 (35)	8.3 (34)	26,378	3.0 (2)	6.9 (35)	13.7 (36)	9,511	15.1 (16)	16.6 (33)	15,779	7.0 (1)
Inyo	118	10.2 (60)	7.6 (20)	1,387	5.5 (48)	4.3 (7)	14.4 (43)	482	19.1 (60)	13.9 (8)	682	13.9 (30)
Kern	30,418	6.7 (45)	8.0 (28)	131,203	5.1 (38)	5.4 (17)	15.5 (52)	43,658	14.8 (14)	15.9 (22)	61,360	10.9 (7)
Kings	1,814	5.2 (13)	9.8 (51)	14,647	5.4 (44)	4.0 (2)	20.1 (65)	4,977	18.3 (57)	21.4 (62)	9,276	25.9 (64)
L.A. East	61,870	6.0 (27)	6.9 (10)	148,278	6.0 (57)	4.2 (5)	17.7 (60)	31,207	16.6 (41)	17.9 (48)	49,478	15.6 (44)
L.A. North	21,674	6.2 (34)	6.2 (5)	76,989	6.2 (60)	4.9 (14)	16.7 (58)	15,900	15.9 (25)	17.7 (46)	21,891	12.1 (15)
L.A. Other	59,775	7.1 (49)	6.9 (10)	526,166	4.5 (22)	4.2 (5)	18.5 (62)	215,997	16.3 (32)	18.9 (56)	305,949	13.3 (24)
L.A. South	59,735	6.6 (42)	6.6 (8)	142,592	6.6 (61)	4.1 (3)	18.5 (62)	30,636	17.0 (48)	18.2 (50)	48,037	12.6 (18)
L.A. West	6,831	6.5 (38)	6.4 (6)	20,914	7.5 (65)	4.5 (8)	18.6 (64)	5,066	16.7 (42)	16.3 (27)	6,852	13.4 (25)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 - 2008 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Lake	578	7.3 (52)	7.3 (15)	3,219	4.3 (16)	8.3 (53)	12.2 (18)	1,009	14.1 (10)	14.4 (14)	1,707	12.0 (12)
Lassen	497	6.4 (37)	5.0 (3)	2,214	4.8 (33)	10.6 (60)	9.6 (5)	722	12.7 (2)	12.9 (4)	1,013	12.4 (17)
Long Beach	7,979	7.3 (52)	7.5 (18)	34,956	5.3 (43)	4.6 (11)	18.0 (61)	11,749	18.5 (58)	20.5 (61)	13,809	14.1 (33)
Madera	7,750	5.3 (15)	7.4 (17)	27,632	5.1 (38)	5.3 (16)	14.7 (45)	9,473	16.0 (28)	15.5 (20)	18,865	18.1 (52)
Marin	363	4.4 (2)	12.1 (58)	6,783	3.5 (5)	7.2 (39)	12.7 (22)	1,989	16.7 (42)	17.1 (40)	3,327	10.2 (4)
Mariposa	78	* *	* *	395	5.6 (51)	7.8 (48)	11.4 (13)	127	13.4 (6)	14.2 (10)	198	18.7 (58)
Mendocino	2,237	6.3 (35)	7.9 (27)	7,688	3.4 (4)	9.3 (56)	12.3 (20)	2,200	19.3 (61)	19.5 (59)	2,701	17.0 (50)
Merced	10,338	5.9 (24)	8.6 (40)	38,119	5.4 (44)	4.6 (11)	15.9 (53)	12,775	16.7 (42)	17.8 (47)	24,328	18.2 (53)
Modoc	168	4.2 (1)	7.7 (24)	761	4.2 (14)	11.6 (64)	9.9 (6)	249	12.9 (3)	9.2 (2)	352	13.4 (25)
Mono	390	7.4 (54)	5.1 (4)	1,368	5.6 (51)	11.6 (64)	7.5 (1)	356	13.8 (7)	8.4 (1)	762	20.1 (60)
Monterey	5,478	5.1 (10)	9.5 (48)	24,427	5.8 (54)	3.5 (1)	15.4 (49)	5,368	17.7 (54)	18.4 (53)	11,843	13.4 (25)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 - 2008 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Napa	1,441	5.8 (23)	9.6 (49)	10,394	3.5 (5)	10.2 (58)	10.0 (7)	2,613	16.2 (30)	16.9 (37)	3,748	25.7 (63)
Nevada	1,305	5.4 (17)	8.1 (31)	4,091	4.7 (31)	10.8 (62)	8.4 (2)	1,005	17.0 (48)	13.5 (7)	1,521	11.0 (8)
Orange	34,208	5.2 (13)	6.9 (10)	81,124	4.4 (20)	6.1 (28)	13.8 (37)	23,187	15.9 (25)	16.4 (30)	66,733	13.1 (23)
Pasadena	991	7.1 (49)	3.5 (1)	5,985	5.8 (54)	7.2 (39)	14.3 (42)	2,001	16.5 (37)	18.2 (50)	3,216	12.0 (12)
Placer	2,712	6.5 (38)	8.5 (37)	12,793	5.1 (38)	7.4 (43)	11.3 (11)	3,407	14.4 (11)	14.5 (15)	5,345	15.2 (42)
Plumas	352	5.1 (10)	8.0 (28)	1,212	6.1 (59)	10.2 (58)	11.5 (14)	339	13.9 (8)	13.9 (8)	467	8.8 (3)
Riverside	44,973	6.1 (32)	7.1 (13)	128,554	6.0 (57)	5.5 (20)	15.2 (47)	32,118	14.4 (11)	15.2 (17)	48,179	15.0 (40)
Sacramento	14,602	6.0 (27)	9.2 (44)	138,870	4.9 (36)	7.7 (46)	12.5 (21)	43,971	15.5 (19)	15.4 (18)	58,555	18.3 (55)
San Benito	354	6.5 (38)	12.4 (59)	3,214	4.5 (22)	7.3 (41)	16.4 (56)	976	18.8 (59)	23.2 (63)	2,049	14.1 (33)
San Bernar	30,326	6.5 (38)	6.8 (9)	143,891	6.8 (64)	5.5 (20)	15.9 (53)	38,608	15.5 (19)	16.2 (24)	56,584	15.3 (43)
San Diego	54,163	7.5 (57)	8.5 (37)	206,343	4.8 (33)	6.4 (30)	13.6 (35)	61,423	16.3 (32)	16.0 (23)	119,356	12.6 (18)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 - 2008 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
San Franci	2,959	7.4 (54)	6.4 (6)	31,665	4.6 (29)	7.0 (37)	12.2 (18)	9,824	17.2 (51)	16.3 (27)	4,652	12.3 (16)
San Joaqui	22,996	7.4 (54)	8.0 (28)	96,443	5.2 (42)	6.7 (34)	14.6 (44)	31,568	15.4 (18)	16.5 (32)	33,694	14.6 (37)
San Luis O	3,007	5.7 (19)	9.3 (45)	17,455	3.9 (11)	8.2 (51)	10.8 (10)	5,000	15.5 (19)	14.5 (15)	6,995	17.0 (50)
San Mateo	2,431	6.0 (27)	7.5 (18)	32,783	3.6 (7)	7.8 (48)	12.1 (17)	9,337	16.7 (42)	18.0 (49)	13,255	10.8 (5)
Santa Barb	13,626	10.2 (60)	7.3 (15)	58,887	4.6 (29)	6.5 (31)	13.0 (28)	18,271	17.3 (52)	18.3 (52)	34,058	15.8 (46)
Santa Clar	11,521	5.4 (17)	7.6 (20)	99,805	4.5 (22)	6.0 (25)	13.1 (31)	29,115	15.8 (23)	16.6 (33)	39,703	10.8 (5)
Santa Cruz	2,951	5.9 (24)	8.3 (34)	9,800	5.4 (44)	6.0 (25)	15.2 (47)	1,788	18.2 (56)	19.2 (58)	3,103	14.4 (35)
Shasta	5,158	6.6 (42)	9.3 (45)	17,842	5.6 (51)	5.4 (17)	13.2 (33)	5,566	17.6 (53)	15.4 (18)	9,617	12.8 (20)
Sierra	37	* *	* *	150	6.7 (62)	10.7 (61)	11.3 (11)	45	* *	* *	69	* *
Siskiyou	1,081	8.5 (59)	7.6 (20)	4,420	4.8 (33)	9.7 (57)	10.2 (8)	1,396	17.0 (48)	14.2 (10)	2,436	15.6 (44)
Solano	5,753	7.6 (58)	8.8 (41)	27,338	4.5 (22)	7.3 (41)	12.8 (24)	8,549	16.5 (37)	17.4 (43)	9,383	13.4 (25)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 - 2008 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Sonoma	6,055	5.7 (19)	9.7 (50)	28,826	3.6 (7)	7.4 (43)	12.8 (24)	7,905	16.5 (37)	16.2 (24)	7,560	18.4 (57)
Stanislaus	13,748	6.0 (27)	7.8 (25)	59,764	4.2 (14)	6.9 (35)	13.9 (38)	18,924	15.9 (25)	17.1 (40)	37,242	20.5 (61)
Sutter	2,343	5.1 (10)	10.7 (55)	11,447	3.7 (9)	6.5 (31)	12.8 (24)	3,348	14.4 (11)	16.2 (24)	5,637	13.0 (21)
Tehama	1,399	4.4 (2)	10.7 (55)	10,000	3.2 (3)	8.2 (51)	10.2 (8)	2,941	13.3 (5)	13.3 (6)	4,394	14.8 (38)
Trinity	188	4.8 (7)	9.0 (42)	829	4.7 (31)	8.9 (55)	13.0 (28)	264	12.1 (1)	19.7 (60)	453	13.0 (21)
Tulare	15,957	5.9 (24)	7.6 (20)	90,085	5.5 (48)	4.6 (11)	16.5 (57)	30,564	16.1 (29)	16.9 (37)	53,149	18.8 (59)
Tuolumne	357	7.0 (47)	7.8 (25)	1,329	4.1 (13)	7.4 (43)	9.5 (4)	342	18.1 (55)	14.3 (13)	489	16.2 (48)
Ventura	19,486	6.1 (32)	8.2 (32)	74,472	4.5 (22)	5.0 (15)	15.4 (49)	22,834	16.8 (46)	19.0 (57)	47,009	11.1 (9)
Yolo	3,237	4.6 (6)	10.8 (57)	16,926	3.8 (10)	7.7 (46)	13.1 (31)	5,247	15.6 (22)	17.4 (43)	7,563	15.1 (41)
Yuba	1,574	5.3 (15)	13.0 (60)	9,916	4.3 (16)	5.9 (23)	13.3 (34)	3,175	16.3 (32)	16.4 (30)	4,853	15.8 (46)
Unknown Counties	1	* *	* *	5	* *	* *	* *	3	* *	* *	0	* *

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 - 2008 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
California	684,675	6.5	7.6	3,062,887	5.1	5.5	15.5	953,794	16.1	17.2	1,468,958	14.2

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 - 2008 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alameda	49,773	6.8 (33)	2.7 (49)	17.7 (20)	21.9 (27)	26,342	17.5 (56)
Alpine	163	9.2 (63)	1.2 (2)	17.2 (9)	23.9 (45)	132	13.6 (34)
Amador	783	6.4 (25)	2.9 (54)	18.0 (26)	20.2 (12)	507	13.8 (36)
Butte	7,933	7.7 (49)	2.0 (17)	18.7 (40)	21.2 (20)	7,534	12.3 (22)
Calaveras	2,034	5.9 (15)	2.2 (24)	19.5 (52)	24.7 (51)	1,840	13.6 (34)
City Of Be	137	5.1 (6)	0.7 (1)	16.1 (3)	24.8 (53)	128	24.2 (64)
Colusa	1,812	4.8 (4)	2.4 (34)	17.6 (14)	25.7 (62)	1,632	14.9 (43)
Contra Cos	18,375	6.6 (28)	2.1 (21)	18.5 (36)	25.3 (58)	8,034	17.4 (55)
Del Norte	865	6.7 (30)	1.5 (7)	20.9 (61)	22.8 (34)	752	8.4 (3)
El Dorado	2,646	6.8 (33)	2.8 (52)	16.1 (3)	17.6 (4)	1,858	9.6 (8)
Fresno	107,619	8.0 (54)	2.3 (29)	18.1 (29)	24.5 (48)	95,138	14.3 (39)
Glenn	1,182	7.7 (49)	2.6 (43)	18.0 (26)	27.4 (65)	1,188	14.1 (38)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 - 2008 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Humboldt	2,785	5.9 (15)	2.0 (17)	19.7 (54)	21.7 (25)	2,110	13.5 (33)
Imperial	9,137	4.3 (2)	3.0 (58)	17.9 (23)	25.0 (55)	8,961	9.7 (9)
Inyo	671	6.1 (17)	2.4 (34)	17.6 (14)	24.3 (47)	408	10.5 (13)
Kern	54,203	5.3 (8)	3.0 (58)	17.6 (14)	23.7 (43)	43,483	12.0 (20)
Kings	4,904	6.1 (17)	2.4 (34)	18.1 (29)	25.6 (61)	4,740	21.6 (63)
L.A. East	66,924	6.2 (20)	2.4 (34)	17.6 (14)	20.6 (16)	81,521	10.4 (11)
L.A. North	43,876	9.6 (64)	2.2 (24)	19.1 (49)	20.5 (14)	42,855	8.7 (4)
L.A. Other	304,135	4.9 (5)	2.3 (29)	18.8 (41)	25.8 (63)	249,151	11.9 (19)
L.A. South	88,518	8.8 (60)	2.1 (21)	19.0 (47)	18.5 (5)	89,182	8.9 (5)
L.A. West	9,553	5.6 (9)	2.6 (43)	18.0 (26)	21.5 (23)	8,399	14.5 (41)
Lake	1,179	6.8 (33)	1.7 (12)	18.9 (46)	20.9 (18)	928	12.5 (25)
Lassen	1,163	11.7 (65)	1.5 (7)	17.6 (14)	25.1 (57)	932	10.6 (14)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 - 2008 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Long Beach	17,881	5.6 (9)	2.0 (17)	18.8 (41)	24.7 (51)	13,458	15.4 (48)
Madera	11,781	6.9 (39)	2.2 (24)	18.8 (41)	23.1 (37)	12,165	13.0 (29)
Marin	1,916	8.4 (59)	2.0 (17)	19.4 (50)	23.3 (39)	1,020	11.2 (15)
Mariposa	196	8.2 (58)	2.6 (43)	17.3 (11)	16.8 (2)	155	11.6 (17)
Mendocino	2,076	6.4 (25)	1.5 (7)	21.4 (63)	23.6 (42)	1,460	12.7 (27)
Merced	13,988	9.0 (61)	2.2 (24)	18.4 (35)	22.3 (31)	13,444	17.3 (54)
Modoc	345	2.9 (1)	2.6 (43)	19.4 (50)	20.6 (16)	324	7.4 (1)
Mono	377	6.4 (25)	4.0 (64)	17.0 (7)	19.1 (8)	346	19.9 (62)
Monterey	7,327	7.7 (49)	1.3 (3)	20.6 (60)	23.1 (37)	7,472	10.4 (11)
Napa	3,135	6.8 (33)	1.4 (5)	20.0 (57)	25.0 (55)	2,088	19.8 (61)
Nevada	989	7.3 (44)	2.6 (43)	21.2 (62)	19.1 (8)	738	12.6 (26)
Orange	48,906	6.9 (39)	2.6 (43)	18.5 (36)	21.1 (19)	72,899	13.1 (30)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 - 2008 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Pasadena	3,190	5.2 (7)	2.4 (34)	17.9 (23)	23.3 (39)	2,947	7.7 (2)
Placer	3,164	7.7 (49)	3.1 (62)	18.5 (36)	19.4 (10)	2,020	14.9 (43)
Plumas	436	7.6 (48)	2.8 (52)	15.8 (1)	22.0 (29)	285	10.2 (10)
Riverside	47,261	5.7 (12)	2.9 (54)	17.8 (21)	21.8 (26)	43,847	12.8 (28)
Sacramento	68,571	8.0 (54)	2.7 (49)	17.3 (11)	21.5 (23)	46,489	19.6 (60)
San Benito	1,192	9.1 (62)	1.4 (5)	21.8 (64)	26.8 (64)	1,137	16.4 (52)
San Bernar	63,961	6.3 (23)	2.9 (54)	17.8 (21)	21.9 (27)	59,685	13.3 (32)
San Diego	85,084	6.7 (30)	2.3 (29)	18.8 (41)	22.9 (35)	84,034	11.3 (16)
San Franci	19,815	7.7 (49)	3.0 (58)	16.7 (5)	18.6 (6)	6,787	16.1 (50)
San Joaqui	48,589	6.8 (33)	2.9 (54)	17.3 (11)	22.3 (31)	23,390	16.1 (50)
San Luis O	4,736	7.1 (43)	2.4 (34)	19.5 (52)	22.0 (29)	3,669	13.1 (30)
San Mateo	11,133	7.3 (44)	1.7 (12)	20.5 (58)	25.3 (58)	5,411	11.8 (18)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 - 2008 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Santa Barb	15,582	8.0 (54)	1.7 (12)	19.8 (55)	25.4 (60)	14,523	12.4 (23)
Santa Clar	45,015	7.0 (41)	3.0 (58)	17.9 (23)	22.5 (33)	23,399	12.0 (20)
Santa Cruz	2,719	7.4 (46)	1.3 (3)	20.5 (58)	23.9 (45)	2,412	9.2 (7)
Shasta	4,921	7.0 (41)	1.7 (12)	18.3 (32)	19.5 (11)	3,994	15.0 (45)
Sierra	111	8.1 (57)	6.3 (65)	23.4 (65)	13.5 (1)	98	* *
Siskiyou	1,689	4.4 (3)	2.4 (34)	17.6 (14)	20.3 (13)	1,462	14.0 (37)
Solano	9,124	6.2 (20)	2.2 (24)	18.6 (39)	24.5 (48)	4,424	17.0 (53)
Sonoma	7,698	6.7 (30)	2.1 (21)	19.8 (55)	22.9 (35)	3,216	15.2 (47)
Stanislaus	23,844	5.6 (9)	2.3 (29)	18.2 (31)	23.8 (44)	24,064	18.0 (58)
Sutter	2,478	5.7 (12)	3.2 (63)	16.7 (5)	20.5 (14)	2,172	18.3 (59)
Tehama	2,536	5.7 (12)	1.8 (16)	17.2 (9)	21.2 (20)	2,086	14.5 (41)
Trinity	396	6.8 (33)	1.5 (7)	15.9 (2)	18.7 (7)	341	15.0 (45)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2006 - 2008 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Tulare	39,665	6.3 (23)	2.5 (42)	18.3 (32)	24.5 (48)	38,273	15.7 (49)
Tuolumne	437	6.6 (28)	1.6 (11)	18.8 (41)	17.2 (3)	364	12.4 (23)
Ventura	19,833	6.2 (20)	2.3 (29)	19.0 (47)	23.5 (41)	19,524	9.0 (6)
Yolo	5,888	6.1 (17)	2.4 (34)	17.0 (7)	24.9 (54)	3,374	17.8 (57)
Yuba	2,591	7.4 (46)	2.7 (49)	18.3 (32)	21.2 (20)	2,083	14.3 (39)
Unknown Counties	5	* *	* *	* *	* *	2	* *
California	1,430,951	6.6	2.4	18.4	23.0	1,228,836	12.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.