

Table 18C
2010 Pediatric Nutrition Surveillance ⁽¹⁾
California

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years ⁽²⁾

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity ⁽³⁾													
Race/Ethnicity and Year		Birthweight ⁽⁴⁾			Height and Weight ^(5,6)							Anemia Low Hb/Hct ⁽⁷⁾	
		Low	High		Short Stature	Under- weight	Obese	≥ 2 Yrs Overweight and Obese					
		Number	% <2500 g	% >4000 g				Number	%	%	%	Number	% Overweight
White, Not Hispanic	2010	22,193	5.8	8.7	94,134	5.4	4.9	11.2	28,926	15.6	13.8	46,325	13.6
	2009	22,580	6.4	8.9	97,178	5.5	5.0	11.6	29,528	14.9	13.7	43,347	13.6
	2008	21,349	6.9	9.0	86,119	5.6	4.8	11.4	24,990	15.0	13.6	38,419	13.4
	2007	20,693	7.1	8.8	85,430	5.7	4.7	10.9	24,269	15.2	13.7	36,773	13.2
	2006	21,767	6.6	8.9	89,232	5.6	4.9	10.5	26,160	15.8	13.0	39,145	12.8
	2005	17,202	7.4	9.1	74,990	5.5	4.8	11.1	26,238	15.5	13.4	37,153	12.1
	2004	17,570	8.2	9.8	79,204	5.5	4.8	11.6	28,066	15.6	13.5	39,334	12.1
	2003	16,347	7.8	10.2	78,415	5.1	5.0	11.3	27,919	15.3	13.1	37,832	12.2
	2002	15,825	7.0	10.4	80,207	5.2	4.8	11.5	28,743	15.3	13.8	38,533	11.7
	2001	13,214	6.8	10.7	74,618	5.5	5.1	10.8	26,702	15.5	12.3	35,973	12.6
Black, Not Hispanic	2010	9,551	10.4	5.3	44,625	7.6	5.1	12.9	14,622	14.8	13.2	18,443	19.9
	2009	10,484	11.1	4.7	49,042	7.5	5.1	13.3	15,648	14.6	13.2	19,066	20.3
	2008	10,675	11.6	5.1	45,778	7.7	5.1	12.9	14,005	14.9	13.2	17,826	19.7
	2007	10,329	11.6	5.2	48,748	7.7	4.7	13.4	15,435	14.4	14.1	19,282	19.2
	2006	10,670	11.8	5.3	52,552	7.3	4.6	13.1	17,869	14.5	13.4	22,049	18.2
	2005	8,227	11.9	6.0	45,545	7.1	4.7	13.2	18,290	14.8	13.2	21,494	16.5
	2004	8,651	12.3	5.9	47,749	6.9	4.9	13.5	19,570	14.2	13.3	23,837	16.8
	2003	8,401	12.1	5.4	47,747	6.8	4.8	13.1	19,609	14.4	12.9	23,882	17.7
	2002	8,217	12.2	6.3	47,494	7.2	4.7	13.4	19,267	14.3	13.4	22,962	16.5
	2001	6,704	11.6	6.3	44,774	7.3	4.9	13.1	18,073	14.3	12.2	21,801	18.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(6) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 18C
2010 Pediatric Nutrition Surveillance ⁽¹⁾
California

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years ⁽²⁾

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity ⁽³⁾													
Race/Ethnicity and Year		Birthweight ⁽⁴⁾			Height and Weight ^(5,6)							Anemia Low Hb/Hct ⁽⁷⁾	
		Low	High		Short Stature	Under- weight	Obese	≥ 2 Yrs Overweight and Obese					
		Number	% <2500 g	% >4000 g				Number	%	%	%	Number	% Overweight
Hispanic	2010	123,540	5.8	7.3	550,796	5.3	3.5	15.4	185,277	16.6	18.8	304,862	12.9
	2009	146,501	5.7	7.3	666,054	5.2	3.4	15.7	221,562	16.4	18.4	360,791	13.4
	2008	157,826	6.1	7.5	649,602	5.3	3.4	15.3	203,364	16.5	18.5	336,952	13.9
	2007	149,254	5.8	7.6	673,034	5.3	3.3	15.3	211,649	16.8	18.7	345,444	13.8
	2006	159,477	5.9	7.8	716,350	5.1	3.4	15.2	236,597	16.6	18.2	393,813	13.7
	2005	124,020	6.8	8.2	587,987	5.1	3.3	15.9	230,859	16.7	18.8	367,681	13.2
	2004	123,984	7.0	8.4	594,514	5.1	3.3	16.2	237,147	16.7	18.8	378,307	14.0
	2003	115,362	6.6	8.7	591,108	5.0	3.4	16.3	242,274	16.7	19.0	374,196	13.6
	2002	112,084	6.1	9.3	587,094	5.1	3.4	16.2	237,925	16.8	18.6	363,388	13.0
	2001	93,404	6.1	9.4	549,881	5.0	3.3	15.7	221,484	16.7	17.7	340,440	13.4
American Indian/Alaska Native	2010	499	5.8	9.0	2,750	5.2	4.4	17.6	1,015	19.0	22.8	1,418	11.8
	2009	510	4.9	9.2	3,421	4.8	4.2	16.2	1,208	18.0	21.2	1,754	16.9
	2008	539	7.1	10.4	3,096	5.6	4.4	14.6	1,012	17.9	20.6	1,557	14.1
	2007	515	9.3	9.5	2,879	5.4	3.7	14.5	946	19.1	20.2	1,486	14.3
	2006	559	5.4	10.4	2,824	5.5	4.5	16.1	960	18.5	21.4	1,503	13.6
	2005	433	6.2	10.2	2,204	5.6	3.4	16.2	824	20.1	19.4	1,274	14.2
	2004	407	7.6	8.8	2,202	5.4	4.0	16.3	876	17.2	19.7	1,304	12.6
	2003	381	8.4	13.6	2,255	6.7	4.3	16.7	872	15.7	21.8	1,324	11.9
	2002	362	6.1	11.0	2,356	5.3	4.0	17.7	959	19.5	21.4	1,341	12.5
	2001	384	4.7	8.9	2,552	7.1	3.7	16.9	1,007	20.2	19.7	1,489	15.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(6) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 18C
2010 Pediatric Nutrition Surveillance ⁽¹⁾
California

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years ⁽²⁾

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity ⁽³⁾													
Race/Ethnicity and Year		Birthweight ⁽⁴⁾			Height and Weight ^(5,6)							Anemia Low Hb/Hct ⁽⁷⁾	
		Low	High		Short Stature	Under- weight	Obese	≥ 2 Yrs Overweight and Obese					
		Number	% <2500 g	% >4000 g				Number	%	%	%	Number	% Overweight
Asian	2010	4,789	6.7	3.8	24,428	6.1	4.7	10.2	8,356	12.9	11.9	9,360	14.1
	2009	6,168	6.9	4.0	31,637	6.5	4.5	11.3	10,813	13.1	13.4	12,319	15.6
	2008	6,498	7.1	3.9	32,635	7.2	4.0	11.3	10,978	13.3	13.2	11,894	16.0
	2007	6,018	6.3	3.3	34,175	6.8	4.1	11.1	11,782	13.0	12.7	11,572	16.1
	2006	5,685	7.1	4.3	35,605	7.2	4.0	11.1	12,490	13.4	13.2	12,964	16.1
	2005	4,717	7.3	4.7	30,610	7.2	4.2	11.0	12,665	13.8	12.5	13,238	14.6
	2004	5,410	6.8	5.1	31,490	7.3	3.9	12.5	12,431	14.6	15.3	13,468	15.2
	2003	8,736	6.7	6.5	54,395	5.3	4.0	12.1	18,282	14.3	15.2	22,761	14.3
	2002	6,637	7.4	5.4	43,858	6.0	3.8	12.5	14,692	14.5	15.3	19,224	13.5
	2001	5,920	7.7	5.5	34,796	6.5	4.3	11.7	11,996	14.5	14.7	13,620	13.9
Pacific Islander	2010	367	6.0	11.4	2,536	5.2	4.4	16.2	910	17.4	22.0	1,326	12.5
	2009	513	5.7	11.3	2,648	5.0	3.7	16.6	856	18.5	21.3	1,189	11.8
	2008	420	6.2	11.2	1,744	4.5	3.5	18.2	500	20.6	25.4	796	13.2
	2007	344	6.4	10.5	1,696	4.4	3.2	17.3	496	21.4	25.4	754	13.8
	2006	379	7.1	10.3	1,715	4.2	3.4	19.7	550	18.2	25.6	737	13.0
	2005	296	5.1	14.2	1,483	3.4	3.6	21.5	530	21.9	25.8	698	10.2
	2004	0	*	*	0	*	*	*	0	*	*	0	*
	2003	0	*	*	0	*	*	*	0	*	*	0	*
	2002	0	*	*	0	*	*	*	0	*	*	0	*
	2001	0	*	*	0	*	*	*	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(6) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 18C
2010 Pediatric Nutrition Surveillance ⁽¹⁾
California

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years ⁽²⁾

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity ⁽³⁾													
Race/Ethnicity and Year		Birthweight ⁽⁴⁾			Height and Weight ^(5,6)							Anemia Low Hb/Hct ⁽⁷⁾	
		Low	High		Short Stature	Under- weight	Obese	≥ 2 Yrs Overweight and Obese					
		Number	% <2500 g	% >4000 g				Number	%	%	%	Number	% Overweight
Filipino	2010	676	9.3	5.6	3,158	8.2	4.2	11.1	883	11.8	13.7	1,051	12.8
	2009	817	8.1	2.8	4,151	9.2	4.7	11.1	1,271	13.5	13.4	1,309	12.7
	2008	917	8.1	3.4	4,258	7.4	4.6	11.1	1,271	13.8	15.0	1,437	12.9
	2007	882	8.6	3.3	4,415	7.7	5.0	10.4	1,335	13.3	12.3	1,645	12.7
	2006	1,015	10.2	3.9	4,880	8.1	4.4	10.8	1,600	13.1	14.4	1,953	11.1
	2005	885	8.6	4.3	4,323	7.4	4.1	11.4	1,583	13.8	14.7	1,837	10.9
	2004	0	*	*	0	*	*	*	0	*	*	0	*
	2003	0	*	*	0	*	*	*	0	*	*	0	*
	2002	0	*	*	0	*	*	*	0	*	*	0	*
	2001	0	*	*	0	*	*	*	0	*	*	0	*
All Other	2010	30,185	6.7	7.6	162,033	5.6	5.2	11.5	44,517	15.2	15.5	59,176	14.6
	2009	34,623	6.5	7.5	185,533	5.4	5.0	11.7	51,777	15.2	15.2	65,527	15.1
	2008	35,667	7.4	8.5	170,297	5.3	5.1	11.6	45,523	15.2	15.6	57,647	15.0
	2007	32,893	6.8	8.0	170,906	5.3	4.9	11.8	46,278	15.4	15.9	57,606	15.3
	2006	30,304	6.7	8.0	158,198	5.5	4.8	11.7	43,735	15.9	15.3	55,704	14.8
	2005	22,385	7.7	8.4	121,025	5.4	4.7	12.1	40,986	16.0	15.5	49,911	13.9
	2004	13,568	8.2	8.3	73,645	5.9	4.8	12.2	24,483	15.7	15.6	31,017	14.5
	2003	11,270	7.8	8.2	62,570	5.8	5.1	11.8	20,686	15.0	15.0	25,425	14.0
	2002	9,199	7.9	8.5	53,964	5.9	5.2	12.3	17,768	14.9	15.4	22,270	14.1
	2001	7,606	7.5	8.4	44,547	6.1	4.9	12.4	14,512	15.3	14.8	18,771	15.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(6) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 18C
2010 Pediatric Nutrition Surveillance (1)
California

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity (3)													
Race/Ethnicity and Year		Birthweight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
		Low	High		Short Stature	Under- weight	Obese	≥ 2 Yrs Overweight and Obese					
		Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese		
Total All Race/Ethnic Groups	2010	191,800	6.2	7.3	884,460	5.5	4.1	14.0	284,506	16.1	17.3	441,961	13.5
	2009	222,196	6.2	7.3	1,039,664	5.4	4.0	14.3	332,663	15.9	17.0	505,302	14.0
	2008	233,891	6.7	7.6	993,529	5.5	3.9	14.1	301,643	16.0	17.3	466,528	14.3
	2007	220,928	6.4	7.5	1,021,283	5.5	3.8	14.1	312,190	16.2	17.4	474,562	14.2
	2006	229,856	6.4	7.7	1,061,356	5.4	3.8	14.0	339,961	16.2	17.0	527,868	14.0
	2005	178,165	7.2	8.1	868,167	5.4	3.7	14.6	331,975	16.3	17.4	493,286	13.4
	2004	169,590	7.4	8.3	828,804	5.4	3.7	15.1	322,573	16.3	17.6	487,267	14.0
	2003	160,497	7.1	8.6	836,490	5.2	3.8	15.1	329,642	16.2	17.7	485,420	13.8
	2002	152,324	6.7	9.0	814,973	5.3	3.8	15.1	319,354	16.3	17.5	467,718	13.1
	2001	127,232	6.6	9.1	751,168	5.3	3.8	14.7	293,774	16.3	16.6	432,094	13.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(6) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 18C
2010 Pediatric Nutrition Surveillance ⁽¹⁾
California

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged 5 to <20 Years ⁽²⁾

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity ⁽³⁾								
Race/Ethnicity and Year		Height and Weight					Anemia Low Hb/Hct (6)	
		Short Stature (4)	Under- weight (5)	Over- weight (5)				
		Number	% <5th	% <5th	% 85th- <95th	% ≥ 95th		
White, Not Hispanic	2010	39,663	5.4	2.9	17.6	20.3	34,823	12.5
	2009	41,112	5.7	2.9	17.2	19.9	31,726	12.9
	2008	34,951	5.6	2.8	17.6	20.1	26,522	12.7
	2007	33,777	5.9	2.7	17.0	20.0	25,359	12.7
	2006	35,696	5.8	2.8	17.2	19.2	27,675	12.8
	2005	37,470	6.3	3.1	17.2	19.4	27,756	12.8
	2004	40,186	6.0	3.1	17.0	18.7	30,027	12.3
	2003	44,445	5.7	3.1	17.3	18.2	33,380	12.8
	2002	46,664	6.0	3.1	16.7	17.7	34,688	12.4
	2001	42,827	5.9	3.2	16.7	16.8	31,892	12.3
Black, Not Hispanic	2010	27,903	3.7	2.7	17.8	21.1	20,744	24.6
	2009	31,064	4.0	2.6	17.6	21.2	21,577	24.5
	2008	29,079	4.0	2.4	17.7	21.2	20,523	23.7
	2007	32,101	4.2	2.4	17.9	21.2	23,723	23.6
	2006	35,772	4.1	2.5	17.3	21.1	27,331	23.2
	2005	37,337	4.1	2.5	17.7	20.7	27,644	21.7
	2004	40,157	3.9	2.5	17.8	20.2	30,593	21.5
	2003	40,349	4.3	2.7	16.8	19.3	30,544	22.3
	2002	38,428	4.1	2.6	17.2	18.8	28,859	21.0
	2001	35,294	4.0	2.8	16.5	17.5	26,428	21.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for height-for-age.

(5) Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(6) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 18C
2010 Pediatric Nutrition Surveillance (1)
California

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged 5 to <20 Years (2)

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity (3)								
Race/Ethnicity and Year		Height and Weight					Anemia Low Hb/Hct (6)	
		Short Stature (4)	Under- weight (5)	Over- weight (5)				
		Number	% <5th	% <5th	% 85th- <95th	% ≥ 95th		
Hispanic	2010	316,261	6.1	2.3	19.4	24.7	321,855	10.4
	2009	355,675	6.3	2.2	19.1	24.5	339,165	10.9
	2008	313,161	6.4	2.1	18.9	24.4	297,374	11.0
	2007	312,265	6.4	2.1	19.1	24.7	293,740	11.4
	2006	320,961	6.1	2.2	19.2	24.8	308,834	11.5
	2005	322,941	6.5	2.2	19.0	24.1	308,698	11.1
	2004	330,393	6.2	2.3	18.9	23.8	312,758	11.5
	2003	337,291	6.1	2.3	19.4	23.2	331,214	11.5
	2002	342,337	6.5	2.4	18.8	22.2	326,786	10.8
	2001	326,131	6.7	2.4	18.5	21.2	304,949	11.4
American Indian/Alaska Native	2010	1,214	4.6	1.5	18.1	27.5	903	12.2
	2009	1,503	5.3	2.0	19.9	29.5	1,182	13.0
	2008	1,395	5.5	2.2	20.1	26.2	1,076	13.9
	2007	1,311	5.3	1.8	21.2	29.2	1,104	12.4
	2006	1,272	5.0	2.6	19.7	27.2	1,035	12.5
	2005	1,144	6.0	1.5	21.2	28.8	962	12.7
	2004	1,283	5.4	1.9	20.3	26.2	1,038	12.5
	2003	1,340	6.9	2.5	18.3	27.0	1,141	10.4
	2002	1,455	4.9	2.1	20.5	24.8	1,239	11.3
	2001	1,406	6.0	2.3	18.7	24.7	1,154	13.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for height-for-age.

(5) Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(6) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 18C
2010 Pediatric Nutrition Surveillance ⁽¹⁾
California

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged 5 to <20 Years ⁽²⁾

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity ⁽³⁾								
Race/Ethnicity and Year		Height and Weight					Anemia Low Hb/Hct (6)	
		Short Stature (4)	Under- weight (5)	Over- weight (5)				
		Number	% <5th	% <5th	% 85th- <95th	% ≥ 95th		
Asian	2010	19,546	10.2	5.2	13.9	12.6	13,465	13.7
	2009	26,709	11.6	5.1	13.8	13.3	17,880	15.6
	2008	27,382	13.2	4.7	14.1	13.5	16,589	16.3
	2007	29,045	12.8	4.7	13.8	13.5	15,956	16.7
	2006	29,719	13.5	4.9	13.5	13.8	17,828	15.5
	2005	31,420	14.8	5.2	13.3	13.4	18,514	15.4
	2004	27,600	13.8	5.0	14.1	15.2	17,252	14.1
	2003	36,958	11.4	4.8	14.0	15.2	20,517	14.4
	2002	32,276	12.3	5.0	13.8	14.2	19,110	14.0
	2001	27,026	12.6	5.1	13.7	13.2	16,326	15.1
Pacific Islander	2010	1,393	5.3	3.2	19.6	30.8	1,227	11.7
	2009	1,412	6.7	2.7	17.7	31.1	1,112	12.1
	2008	913	4.4	1.8	21.0	35.5	745	10.7
	2007	809	4.7	1.5	21.5	38.3	632	16.1
	2006	884	4.4	1.6	19.7	38.0	755	14.3
	2005	885	3.6	2.5	21.2	36.6	728	11.4
	2004	0	*	*	*	*	0	*
	2003	0	*	*	*	*	0	*
	2002	0	*	*	*	*	0	*
	2001	0	*	*	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for height-for-age.

(5) Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(6) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 18C
2010 Pediatric Nutrition Surveillance (1)
California

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged 5 to <20 Years (2)

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity (3)								
Race/Ethnicity and Year		Height and Weight					Anemia Low Hb/Hct (6)	
		Short Stature (4)	Under- weight (5)	Over- weight (5)				
		Number	% <5th	% <5th	% 85th- <95th	% ≥ 95th		
Filipino	2010	2,220	10.8	4.5	16.4	19.2	1,650	10.1
	2009	2,748	10.2	4.8	16.6	18.8	1,884	11.3
	2008	2,672	11.5	4.7	14.6	18.9	1,926	11.4
	2007	2,793	11.0	4.8	14.6	18.9	2,066	10.8
	2006	2,722	9.8	4.7	15.4	18.0	2,076	10.9
	2005	2,675	10.7	3.7	15.9	20.0	2,076	12.0
	2004	0	*	*	*	*	0	*
	2003	0	*	*	*	*	0	*
	2002	0	*	*	*	*	0	*
	2001	0	*	*	*	*	0	*
All Other	2010	60,720	6.5	3.3	17.7	22.0	43,299	14.9
	2009	71,155	6.7	3.1	17.8	21.9	45,485	16.0
	2008	61,902	6.7	3.0	17.9	21.6	39,128	16.1
	2007	61,083	6.4	2.8	18.2	22.4	37,487	16.6
	2006	59,286	6.8	2.7	18.1	22.0	39,352	16.3
	2005	56,808	6.7	2.9	17.6	22.4	36,612	15.2
	2004	34,008	6.9	3.1	17.8	20.9	23,571	15.8
	2003	29,631	6.8	3.4	16.9	20.8	20,826	15.2
	2002	26,326	7.3	3.5	17.4	19.3	19,020	14.6
	2001	21,665	7.2	3.4	16.3	18.3	16,506	15.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for height-for-age.

(5) Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(6) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 18C
2010 Pediatric Nutrition Surveillance (1)
California

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged 5 to <20 Years (2)

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity (3)								
Race/Ethnicity and Year		Height and Weight					Anemia Low Hb/Hct (6)	
		Short Stature (4)	Under- weight (5)	Over- weight (5)				
		Number	% <5th	% <5th	% 85th- <95th	% ≥ 95th		
Total All Race/Ethnic Groups	2010	468,920	6.1	2.6	18.7	23.3	437,966	11.8
	2009	531,378	6.5	2.5	18.4	23.1	460,011	12.4
	2008	471,455	6.6	2.4	18.3	22.8	403,883	12.5
	2007	473,184	6.6	2.4	18.4	23.1	400,067	12.9
	2006	486,312	6.5	2.5	18.4	23.1	424,886	12.9
	2005	490,680	6.9	2.5	18.3	22.7	422,990	12.5
	2004	473,627	6.5	2.6	18.3	22.4	415,239	12.7
	2003	490,014	6.4	2.7	18.4	21.7	437,622	12.7
	2002	487,486	6.7	2.7	18.1	20.8	429,702	12.0
	2001	454,349	6.8	2.7	17.8	19.9	397,255	12.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for height-for-age.

(5) Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(6) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

