

Table 2C
2010 Pediatric Nutrition Surveillance (1)
California
Summary of Health Indicators
Children Aged < 5 Years (2)

Summary of Health Indicators (3)				
	California Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Birthweight (4)				
Low	191,800	6.2	1,961,342	8.9
High	191,800	7.3	1,961,342	6.4
Height and Weight (5,6)				
< 5 Yrs Short Stature	884,460	5.5	8,317,427	6.4
< 5 Yrs Underweight	884,460	4.1	8,317,427	3.5
< 5 Yrs Obese	884,460	14.0	8,317,427	12.9
≥ 2 Yrs Overweight	284,506	16.1	3,612,436	16.3
≥ 2 Yrs Obese	284,506	17.3	3,612,436	14.7
Anemia (7)				
Low Hb	436,199	13.6	5,201,886	15.1
Low Hct	18,155	8.6	538,293	11.6
Low Hb/Hct	441,961	13.5	5,524,196	14.9
Breastfeeding (8)				
Ever Breastfed	0	*	1,769,160	61.7
Breastfed At Least 6 Months	0	*	911,779	27.0
Breastfed At Least 12 Months	0	*	1,092,573	18.5
Smoking in Household				
≤ 2 Hours TV Viewing/ Day (9)	1,129,707	3.0	4,355,723	10.2
	0	*	1,022,933	77.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis. Low birthweight is defined as < 2500 grams and high birthweight is defined as > 4000 grams.

(5) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(6) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

(8) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(9) Children aged 2 years and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 2C
2010 Pediatric Nutrition Surveillance (1)
California
Summary of Health Indicators
Children Aged 5 to <20 Years (2)

Summary of Health Indicators (3)				
	California Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Height and Weight				
Short Stature (4)	< 5th	465,332 6.2	0	*
Underweight (5)	< 5th	465,332 2.5	0	*
Overweight (5)	85th - < 95th	465,332 18.8	0	*
	≥ 95th	465,332 23.3	0	*
Anemia (6)				
Low Hb		432,752 11.8	0	*
Low Hct		16,428 10.0	0	*
Low Hb/Hct		437,970 11.8	0	*
Smoking in Household		650,665 3.5	0	*
≤2 Hours TV Viewing/Day (7)		0 *	0	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 2000 CDC growth chart percentiles for height-for-age.

(5) Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - <95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(6) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

(7) Children aged 2 years and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

