

Table 6B
2008 - 2010 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			(8)	
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Alameda	14,718	6.1 (37)	8.9 (40)	84,047	4.6 (24)	4.2 (28)	11.8 (24)	26,601	16.2 (37)	16.6 (33)	19,177	13.5 (31)
Alpine	53	* *	* *	259	5.8 (47)	5.8 (52)	13.5 (42)	95	* *	* *	134	14.9 (46)
Amador	187	3.7 (5)	7.0 (16)	2,343	5.5 (44)	3.7 (18)	12.7 (35)	767	18.1 (58)	16.7 (35)	448	13.4 (29)
Butte	7,045	5.2 (13)	8.6 (38)	27,648	4.2 (11)	4.1 (24)	11.0 (20)	9,600	15.6 (21)	13.8 (11)	17,030	12.7 (21)
Calaveras	632	5.4 (19)	8.9 (40)	4,659	4.6 (24)	4.0 (22)	13.9 (45)	1,786	15.9 (29)	18.4 (54)	2,471	12.0 (15)
City Of Be	81	* *	* *	331	6.6 (57)	5.1 (45)	8.8 (8)	84	* *	* *	151	16.6 (53)
Colusa	910	5.4 (19)	10.1 (53)	4,549	3.7 (4)	4.8 (41)	12.9 (39)	1,381	15.1 (13)	15.3 (19)	2,027	20.1 (63)
Contra Cos	8,874	6.2 (41)	9.7 (50)	41,311	4.2 (11)	5.6 (48)	10.8 (19)	10,885	15.8 (25)	15.4 (21)	8,133	14.0 (36)
Del Norte	399	6.0 (33)	12.5 (62)	2,208	5.0 (35)	2.9 (2)	15.4 (58)	947	19.4 (63)	16.7 (35)	1,543	6.7 (2)
El Dorado	2,111	5.3 (15)	7.0 (16)	9,341	4.4 (17)	7.0 (59)	8.4 (5)	2,886	14.3 (4)	12.6 (7)	4,229	13.1 (26)
Fresno	32,478	7.2 (56)	6.8 (12)	178,868	6.2 (53)	3.3 (8)	15.1 (57)	61,950	16.5 (42)	17.7 (52)	100,966	14.6 (43)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			(8)	
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Glenn	847	5.0 (10)	9.7 (50)	5,371	4.5 (21)	3.0 (4)	14.9 (56)	1,742	16.6 (44)	17.3 (45)	3,508	19.4 (61)
Humboldt	1,892	5.8 (26)	9.5 (47)	13,006	5.4 (42)	3.1 (5)	12.4 (33)	3,960	17.0 (48)	16.7 (35)	6,062	11.7 (13)
Imperial	8,048	6.0 (33)	8.8 (39)	26,660	3.1 (1)	4.2 (28)	13.1 (40)	9,771	14.4 (7)	15.2 (17)	17,058	6.2 (1)
Inyo	93	* *	* *	1,460	6.0 (50)	3.5 (14)	14.0 (46)	526	17.1 (52)	15.2 (17)	274	14.6 (43)
Kern	28,066	6.8 (52)	7.2 (20)	133,126	5.3 (41)	4.0 (22)	14.6 (53)	46,211	15.1 (13)	16.0 (27)	45,247	13.0 (23)
Kings	2,848	6.2 (41)	9.6 (48)	15,745	6.2 (53)	3.1 (5)	17.9 (66)	5,270	17.6 (56)	21.1 (62)	9,896	23.3 (64)
L.A. East	56,238	6.0 (33)	6.8 (12)	139,555	6.7 (59)	3.2 (7)	15.5 (59)	27,631	16.1 (35)	17.6 (48)	40,414	14.8 (45)
L.A. North	22,115	6.2 (41)	5.2 (5)	71,762	7.3 (62)	4.1 (24)	14.7 (54)	13,130	15.5 (20)	18.7 (56)	16,275	12.6 (19)
L.A. Other	57,383	7.0 (54)	6.6 (10)	511,953	4.9 (32)	2.9 (2)	17.2 (65)	211,245	16.3 (38)	19.0 (57)	288,905	13.2 (28)
L.A. South	58,438	6.8 (52)	6.3 (8)	135,974	7.6 (64)	3.4 (10)	15.9 (62)	26,690	16.4 (41)	18.0 (53)	37,187	14.0 (36)
L.A. West	4,801	6.5 (49)	6.3 (8)	13,386	8.0 (65)	3.4 (10)	16.4 (64)	3,022	17.0 (48)	17.6 (48)	3,775	11.4 (12)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			(8)	
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Lake	467	3.6 (4)	8.4 (37)	3,399	4.8 (30)	5.1 (45)	12.0 (28)	1,192	14.3 (4)	14.7 (15)	1,841	13.4 (29)
Lassen	498	7.6 (58)	4.6 (3)	2,027	4.8 (30)	8.2 (63)	9.4 (9)	621	14.3 (4)	12.4 (4)	712	9.7 (3)
Long Beach	10,423	7.7 (61)	7.7 (29)	32,164	6.5 (56)	3.3 (8)	16.2 (63)	8,640	18.3 (59)	21.0 (61)	10,599	15.2 (48)
Madera	7,915	6.1 (37)	7.4 (25)	29,695	5.0 (35)	3.4 (10)	13.5 (42)	11,118	15.9 (29)	15.5 (23)	20,865	16.1 (52)
Marin	1,044	6.2 (41)	8.0 (34)	8,156	3.7 (4)	3.9 (21)	10.7 (17)	2,266	15.4 (18)	14.3 (13)	4,279	9.7 (3)
Mariposa	111	5.4 (19)	5.4 (6)	539	6.1 (52)	9.5 (66)	10.4 (14)	162	19.1 (62)	16.7 (35)	262	10.3 (6)
Mendocino	1,954	5.2 (13)	7.8 (32)	8,322	3.5 (2)	5.6 (48)	12.2 (31)	2,583	18.4 (61)	20.4 (60)	2,397	17.3 (55)
Merced	7,723	6.5 (49)	7.3 (23)	32,217	6.2 (53)	3.7 (18)	14.7 (54)	9,411	16.7 (46)	18.6 (55)	19,579	18.8 (59)
Modoc	155	2.6 (1)	5.8 (7)	836	4.4 (17)	6.9 (58)	9.6 (11)	294	15.6 (21)	12.2 (3)	317	10.4 (7)
Mono	340	6.5 (49)	3.5 (1)	1,280	3.7 (4)	8.6 (65)	5.6 (1)	343	12.2 (1)	5.8 (1)	628	15.4 (49)
Monterey	6,671	5.1 (12)	9.0 (42)	28,228	7.1 (61)	2.4 (1)	14.0 (46)	4,968	16.6 (44)	19.0 (57)	13,708	13.0 (23)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			(8)	
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Napa	1,155	4.2 (6)	10.7 (56)	9,810	3.5 (2)	5.8 (52)	10.5 (16)	2,734	17.0 (48)	17.2 (43)	4,430	26.3 (65)
Nevada	1,499	4.3 (7)	7.7 (29)	5,209	4.2 (11)	6.8 (57)	6.4 (2)	1,427	13.9 (3)	10.9 (2)	2,165	12.7 (21)
Orange	38,631	5.7 (22)	6.9 (14)	69,934	5.0 (35)	5.3 (47)	12.8 (38)	19,324	16.0 (32)	16.8 (39)	118,620	12.5 (18)
Pasadena	1,258	6.1 (37)	4.1 (2)	4,641	4.5 (21)	7.3 (61)	9.4 (9)	1,242	17.1 (52)	16.8 (39)	1,679	13.5 (31)
Placer	2,368	5.0 (10)	7.4 (25)	13,924	4.7 (28)	5.0 (44)	10.3 (13)	4,175	13.8 (2)	12.5 (5)	5,690	11.8 (14)
Plumas	433	5.3 (15)	5.1 (4)	1,536	5.7 (45)	7.2 (60)	8.7 (6)	436	14.4 (7)	12.8 (8)	520	12.3 (17)
Riverside	49,590	5.8 (26)	7.2 (20)	117,095	6.7 (59)	4.2 (28)	14.0 (46)	28,639	14.5 (9)	15.5 (23)	54,766	13.5 (31)
Sacramento	12,985	5.7 (22)	9.1 (43)	132,290	4.9 (32)	5.9 (54)	10.7 (17)	43,536	14.9 (12)	14.6 (14)	53,102	18.9 (60)
San Benito	197	7.6 (58)	9.1 (43)	3,406	4.3 (15)	3.8 (20)	15.5 (59)	1,142	18.0 (57)	23.1 (63)	2,076	13.0 (23)
San Bernar	29,551	6.2 (41)	6.6 (10)	115,582	7.4 (63)	4.5 (34)	14.0 (46)	29,641	15.3 (16)	15.5 (23)	50,542	14.0 (36)
San Diego	45,423	7.6 (58)	9.1 (43)	193,202	5.2 (40)	4.6 (35)	12.4 (33)	59,292	16.1 (35)	16.0 (27)	122,325	13.1 (26)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese			(8)	
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
San Franci	2,641	7.0 (54)	6.9 (14)	31,357	4.7 (28)	4.7 (39)	11.5 (21)	10,418	16.0 (32)	15.5 (23)	4,617	12.2 (16)
San Joaqui	18,899	7.4 (57)	7.6 (28)	96,256	5.4 (42)	4.9 (43)	14.1 (50)	33,179	15.3 (16)	16.9 (41)	30,362	13.9 (35)
San Luis O	2,196	5.3 (15)	10.0 (52)	14,480	4.2 (11)	4.8 (41)	10.4 (14)	4,449	15.1 (13)	14.1 (12)	5,353	17.8 (57)
San Mateo	2,205	6.1 (37)	9.4 (46)	34,428	3.7 (4)	4.6 (35)	11.6 (22)	10,236	17.5 (55)	17.4 (46)	12,593	10.7 (8)
Santa Barb	11,389	6.4 (48)	7.2 (20)	57,012	4.6 (24)	4.1 (24)	11.9 (25)	19,838	17.2 (54)	17.0 (42)	32,992	14.3 (41)
Santa Clar	9,975	5.7 (22)	7.3 (23)	90,055	5.1 (38)	4.2 (28)	11.9 (25)	25,309	16.0 (32)	17.2 (43)	27,529	11.3 (11)
Santa Cruz	2,615	6.3 (46)	7.0 (16)	10,878	5.8 (47)	4.1 (24)	12.7 (35)	1,748	16.5 (42)	16.5 (32)	3,574	14.4 (42)
Shasta	4,419	5.8 (26)	10.9 (58)	17,258	5.9 (49)	3.6 (16)	12.3 (32)	5,603	17.0 (48)	15.3 (19)	8,306	12.6 (19)
Sierra	37	* *	* *	151	9.9 (66)	7.9 (62)	6.6 (3)	46	* *	* *	67	* *
Siskiyou	862	8.6 (62)	8.1 (36)	4,249	4.4 (17)	6.4 (56)	8.7 (6)	1,445	15.6 (21)	12.5 (5)	2,396	17.6 (56)
Solano	3,986	5.9 (29)	10.4 (55)	21,748	4.9 (32)	4.4 (33)	11.7 (23)	7,316	15.8 (25)	16.0 (27)	6,703	11.2 (10)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Sonoma	6,867	5.7 (22)	9.6 (48)	26,353	4.5 (21)	4.7 (39)	12.1 (29)	7,537	15.8 (25)	15.4 (21)	6,221	16.9 (54)
Stanislaus	13,290	6.3 (46)	7.5 (27)	61,763	5.1 (38)	4.2 (28)	13.4 (41)	20,335	16.3 (38)	17.6 (48)	35,556	19.7 (62)
Sutter	1,981	5.9 (29)	10.3 (54)	10,967	4.0 (10)	4.6 (35)	12.1 (29)	3,389	14.5 (9)	16.3 (30)	5,290	10.9 (9)
Tehama	939	5.9 (29)	11.2 (59)	9,145	3.8 (8)	5.7 (50)	9.9 (12)	2,957	14.6 (11)	13.6 (10)	3,621	14.1 (39)
Trinity	150	2.7 (2)	10.7 (56)	662	6.0 (50)	6.2 (55)	13.6 (44)	221	15.4 (18)	16.3 (30)	345	13.6 (34)
Tulare	13,925	6.0 (33)	7.7 (29)	85,456	5.7 (45)	3.5 (14)	15.6 (61)	29,505	15.6 (21)	17.4 (46)	48,087	18.3 (58)
Tuolumne	424	3.3 (3)	8.0 (34)	1,729	3.8 (8)	8.2 (63)	7.8 (4)	481	15.8 (25)	13.5 (9)	379	15.6 (51)
Ventura	16,918	5.9 (29)	7.9 (33)	79,176	4.6 (24)	3.4 (10)	14.3 (52)	26,495	16.3 (38)	19.2 (59)	50,921	10.2 (5)
Yolo	3,839	4.4 (8)	11.2 (59)	16,437	4.3 (15)	4.6 (35)	12.7 (35)	5,442	15.9 (29)	16.6 (33)	7,569	14.1 (39)
Yuba	1,455	4.6 (9)	12.2 (61)	9,959	4.4 (17)	3.6 (16)	11.9 (25)	3,174	16.7 (46)	14.9 (16)	4,934	15.1 (47)
Unknown Counties	227	5.3 (15)	7.0 (16)	1,080	6.6 (57)	5.7 (50)	14.2 (51)	323	18.3 (59)	17.6 (48)	364	15.4 (49)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
County	Birthweight (5)			Height and Weight (3,4)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	%	%	Number	%	%	%	Number	%	%	Number	%
	<2500 g (Rank)	>4000 g (Rank)		(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)
California	647,887	6.4	7.4	2,917,653	5.5	4.0	14.1	918,812	16.0	17.2	1,413,791	13.9

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.
 - (7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators								(3,4)
County	Height and Weight					Anemia Low Hb/Hct (7)		
	Short Stature (5)	Under- weight (6)	Over- weight (6)					
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)	
Alameda	44,675	6.5 (36)	2.8 (53)	17.8 (24)	22.2 (29)	20,630	17.5 (55)	
Alpine	106	7.5 (53)	1.9 (9)	11.3 (1)	27.4 (63)	90	* *	
Amador	761	5.3 (8)	2.5 (40)	18.1 (31)	19.8 (13)	293	14.3 (40)	
Butte	8,304	6.3 (30)	2.3 (29)	19.0 (46)	21.1 (18)	7,899	13.9 (36)	
Calaveras	2,475	5.5 (10)	2.1 (19)	19.6 (52)	23.2 (38)	2,129	9.8 (8)	
City Of Be	183	2.2 (1)	3.3 (62)	15.8 (5)	24.0 (41)	152	22.4 (64)	
Colusa	1,612	5.5 (10)	2.7 (48)	18.5 (38)	25.4 (60)	1,356	15.9 (49)	
Contra Cos	16,103	6.0 (25)	2.3 (29)	18.2 (35)	24.8 (49)	4,622	17.0 (53)	
Del Norte	801	7.0 (43)	2.1 (19)	19.2 (49)	21.7 (23)	760	8.8 (4)	
El Dorado	3,140	7.1 (47)	3.1 (59)	17.5 (16)	18.5 (6)	2,065	12.3 (21)	
Fresno	106,783	7.5 (53)	2.3 (29)	18.2 (35)	25.2 (56)	88,166	13.8 (35)	
Glenn	1,657	7.4 (52)	2.2 (23)	17.9 (26)	25.2 (56)	1,560	17.2 (54)	

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators								(3,4)
County	Height and Weight					Anemia Low Hb/Hct (7)		
	Short Stature (5)	Under- weight (6)	Over- weight (6)					
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)	
Humboldt	2,975	5.8 (19)	1.7 (5)	21.6 (64)	21.9 (25)	2,153	12.7 (29)	
Imperial	9,519	4.0 (3)	2.8 (53)	17.5 (16)	24.6 (47)	9,116	8.4 (3)	
Inyo	743	3.9 (2)	1.9 (9)	19.7 (53)	22.7 (31)	306	13.4 (33)	
Kern	59,073	5.4 (9)	2.7 (48)	17.7 (20)	24.3 (45)	31,992	12.5 (25)	
Kings	5,455	5.7 (16)	2.5 (40)	18.5 (38)	25.3 (58)	5,255	18.9 (62)	
L.A. East	75,340	5.5 (10)	2.1 (19)	15.9 (6)	18.3 (5)	102,862	7.6 (1)	
L.A. North	43,599	10.2 (65)	2.3 (29)	19.3 (50)	20.8 (16)	39,631	11.1 (12)	
L.A. Other	336,619	5.0 (6)	2.5 (40)	19.0 (46)	25.8 (62)	272,349	11.7 (16)	
L.A. South	91,428	9.2 (62)	1.9 (9)	20.8 (60)	18.0 (4)	83,143	9.5 (7)	
L.A. West	6,623	6.0 (25)	2.3 (29)	17.7 (20)	22.0 (26)	5,363	13.5 (34)	
Lake	1,418	5.1 (7)	2.4 (35)	18.0 (30)	19.7 (12)	1,035	12.4 (23)	
Lassen	1,009	6.9 (41)	2.5 (40)	18.3 (37)	18.6 (7)	632	8.9 (5)	

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators								(3,4)
County	Height and Weight					Anemia Low Hb/Hct (7)		
	Short Stature (5)	Under- weight (6)	Over- weight (6)					
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)	
Long Beach	13,630	6.3 (30)	2.4 (35)	19.0 (46)	25.1 (52)	10,174	15.2 (46)	
Madera	14,317	7.0 (43)	1.9 (9)	18.6 (40)	23.6 (39)	13,709	12.2 (19)	
Marin	2,115	6.3 (30)	2.0 (16)	19.8 (54)	24.4 (46)	1,245	12.5 (25)	
Mariposa	240	9.2 (62)	2.5 (40)	15.0 (2)	21.7 (23)	193	12.4 (23)	
Mendocino	2,614	5.9 (23)	1.7 (5)	21.0 (61)	25.1 (52)	1,481	13.9 (36)	
Merced	10,864	8.1 (58)	2.3 (29)	17.9 (26)	22.3 (30)	11,005	17.6 (56)	
Modoc	303	5.6 (13)	1.7 (5)	21.5 (63)	23.8 (40)	237	10.1 (9)	
Mono	394	7.1 (47)	6.3 (65)	16.5 (9)	17.3 (3)	317	17.7 (58)	
Monterey	7,173	7.8 (55)	1.4 (1)	21.2 (62)	24.2 (44)	7,125	10.1 (9)	
Napa	3,150	7.0 (43)	2.1 (19)	20.6 (58)	25.0 (51)	1,934	18.6 (61)	
Nevada	1,513	6.3 (30)	2.4 (35)	18.8 (43)	17.2 (2)	1,173	11.3 (13)	
Orange	48,363	7.2 (49)	2.5 (40)	18.8 (43)	21.0 (17)	128,203	11.7 (16)	

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators								(3,4)
County	Height and Weight					Anemia Low Hb/Hct (7)		
	Short Stature (5)	Under- weight (6)	Over- weight (6)					
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)	
Pasadena	2,036	4.5 (4)	2.0 (16)	18.1 (31)	23.0 (35)	1,447	9.3 (6)	
Placer	3,762	6.8 (40)	3.0 (57)	16.6 (11)	19.0 (9)	2,146	12.3 (21)	
Plumas	561	7.8 (55)	4.6 (64)	15.9 (6)	22.8 (34)	349	11.5 (15)	
Riverside	45,922	5.7 (16)	3.2 (60)	17.8 (24)	21.5 (21)	52,906	12.7 (29)	
Sacramento	68,116	7.2 (49)	3.0 (57)	17.3 (14)	21.1 (18)	46,562	19.4 (63)	
San Benito	1,415	8.2 (61)	1.5 (2)	21.9 (65)	27.7 (64)	1,244	14.5 (42)	
San Bernar	56,690	6.1 (28)	3.4 (63)	17.6 (18)	21.6 (22)	61,703	12.6 (28)	
San Diego	87,605	6.9 (41)	2.5 (40)	18.8 (43)	23.1 (37)	97,603	11.4 (14)	
San Franci	20,441	7.0 (43)	3.2 (60)	16.9 (12)	19.0 (9)	4,822	16.4 (51)	
San Joaqui	52,864	6.4 (34)	2.8 (53)	17.6 (18)	22.7 (31)	24,642	14.4 (41)	
San Luis O	4,327	6.7 (39)	2.2 (23)	18.6 (40)	22.0 (26)	3,375	14.5 (42)	
San Mateo	11,840	6.4 (34)	1.8 (8)	20.7 (59)	25.1 (52)	5,008	11.9 (18)	

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators								(3,4)
County	Height and Weight					Anemia Low Hb/Hct (7)		
	Short Stature (5)	Under- weight (6)	Over- weight (6)					
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)	
Santa Barb	17,105	8.1 (58)	1.6 (3)	20.3 (57)	25.1 (52)	14,595	12.5 (25)	
Santa Clar	41,166	7.2 (49)	2.9 (56)	18.1 (31)	22.7 (31)	17,476	12.2 (19)	
Santa Cruz	2,541	6.5 (36)	1.6 (3)	20.1 (56)	25.3 (58)	2,261	10.4 (11)	
Shasta	4,423	5.9 (23)	1.9 (9)	17.7 (20)	20.3 (14)	3,242	13.3 (32)	
Sierra	76	* *	* *	* *	* *	61	* *	
Siskiyou	1,641	4.8 (5)	2.2 (23)	16.5 (9)	19.2 (11)	1,366	15.7 (48)	
Solano	8,601	5.6 (13)	2.0 (16)	17.9 (26)	24.8 (49)	3,489	15.1 (45)	
Sonoma	7,196	5.7 (16)	1.9 (9)	19.8 (54)	23.0 (35)	2,894	14.9 (44)	
Stanislaus	27,231	5.8 (19)	2.2 (23)	18.6 (40)	24.6 (47)	26,322	16.5 (52)	
Sutter	2,586	5.8 (19)	2.7 (48)	16.9 (12)	21.1 (18)	2,133	15.4 (47)	
Tehama	2,852	5.8 (19)	2.2 (23)	17.3 (14)	20.7 (15)	1,970	12.7 (29)	
Trinity	322	8.1 (58)	1.9 (9)	16.1 (8)	18.6 (7)	293	17.7 (58)	

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6B
2008 - 2010 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators								(3,4)
County	Height and Weight					Anemia Low Hb/Hct (7)		
	Short Stature (5)	Under- weight (6)	Over- weight (6)					
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)	
Tulare	38,804	6.0 (25)	2.4 (35)	18.1 (31)	25.5 (61)	35,233	14.2 (39)	
Tuolumne	490	8.0 (57)	2.7 (48)	15.1 (3)	17.1 (1)	244	16.0 (50)	
Ventura	22,790	6.1 (28)	2.2 (23)	19.3 (50)	24.1 (42)	21,896	8.3 (2)	
Yolo	6,155	5.6 (13)	2.6 (47)	17.9 (26)	24.1 (42)	3,404	17.6 (56)	
Yuba	2,894	6.6 (38)	2.7 (48)	17.7 (20)	22.0 (26)	2,367	13.9 (36)	
Unknown Counties	636	9.4 (64)	2.4 (35)	15.4 (4)	29.6 (65)	456	18.2 (60)	
California	1,468,165	6.4	2.5	18.5	23.1	1,301,864	12.2	

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

