

**Table 6B**  
**1999 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alameda	4,004	7.3 (34)	10.0 (29)	23,538	6.4 (59)	6.0 (30)	15.5 (44)	9,340	16.4 (36)	15.8 (32)	11,121	14.3 (43)
Alpine	74	* *	* *	392	5.4 (43)	6.9 (42)	15.8 (48)	174	20.1 (63)	16.1 (40)	230	12.6 (31)
Amador	189	10.6 (54)	11.1 (41)	1,050	4.0 (13)	6.3 (33)	13.5 (26)	455	15.6 (22)	12.5 (10)	585	11.1 (16)
Butte	1,469	7.6 (36)	12.9 (53)	6,283	4.1 (16)	7.1 (44)	10.6 (6)	2,666	14.8 (11)	10.7 (3)	4,732	9.1 (5)
Calaveras	221	10.4 (52)	9.0 (15)	1,031	5.9 (53)	4.8 (11)	14.7 (38)	446	12.3 (2)	16.8 (46)	588	13.4 (36)
City Of Be	69	* *	* *	455	6.8 (62)	5.9 (27)	14.1 (34)	169	18.9 (60)	17.8 (54)	238	29.4 (64)
Colusa	512	7.0 (31)	9.0 (15)	1,711	4.4 (20)	8.1 (54)	17.6 (59)	585	15.6 (22)	21.4 (63)	946	13.6 (39)
Contra Cos	1,653	8.6 (48)	11.2 (43)	10,735	5.5 (45)	5.9 (27)	15.6 (46)	4,026	18.2 (54)	18.1 (57)	5,156	15.3 (51)
Del Norte	219	4.1 (3)	14.6 (55)	1,050	3.9 (11)	4.8 (11)	14.0 (31)	431	15.1 (13)	15.1 (25)	725	21.7 (62)
El Dorado	720	11.4 (56)	8.5 (9)	3,596	4.7 (29)	7.3 (46)	13.9 (28)	1,387	15.4 (19)	15.9 (34)	2,203	11.6 (20)
Fresno	7,456	6.9 (28)	9.2 (22)	38,695	5.3 (42)	4.1 (3)	15.9 (51)	16,314	15.9 (28)	15.1 (25)	27,109	11.9 (26)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**1999 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Glenn	298	5.4 (8)	14.8 (56)	1,217	3.8 (7)	4.9 (13)	11.4 (9)	483	15.7 (25)	13.9 (15)	953	11.3 (17)
Humboldt	577	4.0 (2)	19.8 (57)	3,885	4.4 (20)	4.9 (13)	13.0 (22)	1,471	18.8 (58)	15.2 (27)	2,304	15.2 (50)
Imperial	937	6.3 (19)	11.8 (47)	5,375	3.1 (1)	7.4 (47)	15.8 (48)	2,142	15.9 (28)	18.0 (56)	3,942	7.6 (2)
Inyo	257	5.4 (8)	11.7 (46)	1,225	5.1 (39)	6.0 (30)	18.4 (62)	487	15.0 (12)	21.1 (62)	734	16.6 (60)
Kern	5,650	6.3 (19)	10.8 (38)	25,145	4.9 (32)	5.4 (24)	16.0 (53)	10,146	16.1 (34)	15.9 (34)	15,181	8.7 (3)
Kings	669	6.9 (28)	9.3 (24)	3,077	4.4 (20)	5.1 (21)	15.5 (44)	1,129	13.8 (7)	14.9 (23)	1,916	19.2 (61)
L.A. East	16,142	6.3 (19)	9.1 (19)	61,148	5.5 (45)	4.1 (3)	17.7 (60)	23,673	16.6 (41)	16.9 (47)	38,962	14.0 (40)
L.A. North	11,389	6.4 (23)	9.1 (19)	55,890	4.9 (32)	5.0 (16)	15.7 (47)	23,292	17.0 (44)	16.9 (47)	36,289	11.5 (19)
L.A. Other	7,404	8.0 (39)	9.9 (28)	73,431	4.4 (20)	4.2 (5)	17.7 (60)	35,602	16.5 (37)	17.0 (49)	50,951	13.5 (38)
L.A. South	17,280	6.9 (28)	9.3 (24)	74,855	5.5 (45)	4.2 (5)	18.5 (63)	32,746	16.5 (37)	17.0 (49)	52,282	11.7 (22)
L.A. West	5,855	7.0 (31)	10.6 (35)	25,626	5.9 (53)	4.5 (8)	17.4 (57)	10,133	16.7 (42)	16.7 (43)	15,881	12.6 (31)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**1999 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Lake	375	8.5 (47)	12.0 (49)	1,503	4.5 (24)	8.6 (57)	12.8 (19)	592	16.0 (32)	14.5 (20)	979	12.5 (29)
Lassen	168	8.9 (49)	10.1 (30)	624	5.8 (52)	8.2 (55)	9.8 (4)	251	19.1 (62)	10.8 (4)	331	11.8 (24)
Long Beach	2,901	8.3 (44)	9.0 (15)	13,375	5.5 (45)	4.4 (7)	17.4 (57)	5,242	17.0 (44)	17.3 (53)	7,552	15.7 (55)
Madera	1,198	6.2 (17)	10.3 (31)	4,943	5.0 (35)	5.3 (23)	14.4 (36)	2,109	15.8 (26)	14.4 (18)	4,244	12.8 (34)
Marin	118	11.0 (55)	9.3 (24)	2,051	3.8 (7)	5.4 (24)	12.0 (16)	761	18.3 (55)	15.4 (28)	1,105	10.1 (10)
Mariposa	44	* *	* *	276	7.2 (64)	7.6 (49)	16.3 (55)	120	17.5 (49)	21.7 (64)	193	10.9 (14)
Mendocino	578	3.5 (1)	11.9 (48)	2,403	3.5 (5)	7.6 (49)	13.5 (26)	942	18.8 (58)	17.0 (49)	1,123	16.1 (58)
Merced	2,247	8.3 (44)	8.7 (12)	9,888	6.4 (59)	3.4 (1)	15.9 (51)	3,917	17.7 (51)	16.7 (43)	7,461	14.5 (46)
Modoc	55	* *	* *	278	5.0 (35)	7.6 (49)	12.9 (21)	116	13.8 (7)	17.2 (52)	181	6.1 (1)
Mono	76	* *	* *	289	6.6 (61)	6.6 (37)	6.9 (1)	102	11.8 (1)	11.8 (7)	150	24.7 (63)
Monterey	1,787	8.2 (43)	10.7 (37)	13,194	3.8 (7)	5.0 (16)	14.9 (41)	5,318	19.0 (61)	18.6 (59)	10,580	12.6 (31)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**1999 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Napa	192	5.2 (7)	12.0 (49)	1,835	3.3 (3)	8.3 (56)	12.5 (17)	593	15.9 (28)	18.4 (58)	723	15.5 (54)
Nevada	274	7.7 (38)	10.9 (39)	1,107	4.5 (24)	9.5 (64)	9.6 (2)	393	17.3 (47)	11.2 (5)	685	10.5 (13)
Orange	11,349	5.7 (11)	9.2 (22)	81,129	4.6 (27)	5.0 (16)	15.8 (48)	36,104	15.8 (26)	15.9 (34)	56,804	15.4 (52)
Pasadena	985	7.1 (33)	4.9 (1)	3,808	4.1 (16)	6.5 (34)	14.0 (31)	1,560	14.2 (9)	15.4 (28)	2,488	8.7 (3)
Placer	384	8.3 (44)	8.9 (14)	2,119	7.0 (63)	7.2 (45)	11.7 (13)	771	15.2 (14)	13.6 (13)	1,218	11.7 (22)
Plumas	103	6.8 (27)	7.8 (5)	670	4.9 (32)	7.5 (48)	13.9 (28)	321	13.4 (5)	15.6 (30)	374	11.0 (15)
Riverside	7,579	6.1 (15)	8.5 (9)	39,737	5.5 (45)	4.7 (9)	16.0 (53)	16,616	15.2 (14)	14.5 (20)	26,838	13.4 (36)
Sacramento	2,856	9.2 (50)	11.6 (45)	27,739	5.0 (35)	6.7 (39)	12.8 (19)	11,858	15.9 (28)	13.4 (12)	17,796	15.1 (48)
San Benito	181	6.1 (15)	7.7 (4)	1,584	6.1 (57)	3.9 (2)	20.2 (64)	659	15.6 (22)	20.6 (61)	1,112	12.8 (34)
San Bernar	9,525	6.5 (24)	8.2 (6)	53,864	6.0 (56)	5.0 (16)	15.3 (43)	23,894	15.3 (17)	14.0 (16)	34,151	15.1 (48)
San Diego	9,988	6.2 (17)	10.5 (33)	61,340	4.6 (27)	6.0 (30)	13.9 (28)	25,966	16.1 (34)	14.8 (22)	40,771	10.1 (10)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**1999 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
San Franci	721	11.4 (56)	7.5 (3)	8,191	6.1 (57)	4.9 (13)	14.8 (40)	3,237	17.7 (51)	16.7 (43)	2,377	9.8 (6)
San Joaqui	3,745	8.1 (42)	8.8 (13)	17,520	5.7 (50)	5.9 (27)	14.9 (41)	7,230	17.5 (49)	15.8 (32)	10,845	11.8 (24)
San Luis O	550	8.0 (39)	10.9 (39)	4,167	4.3 (19)	7.9 (53)	10.4 (5)	1,593	17.3 (47)	12.5 (10)	2,015	12.5 (29)
San Mateo	992	6.6 (25)	9.0 (15)	8,409	4.7 (29)	6.7 (39)	14.0 (31)	3,270	18.5 (57)	19.1 (60)	4,298	10.0 (9)
Santa Barb	2,488	7.6 (36)	8.2 (6)	11,564	4.5 (24)	6.6 (37)	13.4 (24)	4,462	17.0 (44)	16.6 (42)	8,188	11.4 (18)
Santa Clar	2,526	5.0 (6)	10.5 (33)	23,008	4.8 (31)	6.5 (34)	13.0 (22)	8,670	15.5 (20)	15.7 (31)	11,008	15.7 (55)
Santa Cruz	1,080	4.9 (5)	12.6 (52)	7,405	3.9 (11)	5.2 (22)	14.7 (38)	2,881	18.3 (55)	17.9 (55)	4,128	12.1 (27)
Shasta	1,018	9.5 (51)	12.0 (49)	4,303	5.9 (53)	5.8 (26)	11.6 (10)	1,793	14.7 (10)	9.9 (1)	2,863	9.8 (6)
Sierra	6	* *	* *	99	* *	* *	* *	47	* *	* *	55	* *
Siskiyou	242	5.4 (8)	7.4 (2)	1,181	4.0 (13)	8.8 (59)	9.7 (3)	487	17.9 (53)	10.1 (2)	767	14.0 (40)
Solano	1,467	8.0 (39)	8.3 (8)	7,119	5.0 (35)	7.7 (52)	11.9 (15)	2,575	16.7 (42)	13.6 (13)	3,669	14.7 (47)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**1999 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Sonoma	661	6.7 (26)	9.8 (27)	5,804	3.2 (2)	9.0 (61)	10.9 (7)	2,104	16.5 (37)	14.4 (18)	2,691	14.3 (43)
Stanislaus	2,959	10.5 (53)	8.5 (9)	13,934	4.0 (13)	9.1 (62)	12.6 (18)	5,545	15.3 (17)	15.9 (34)	10,585	15.7 (55)
Sutter	425	5.9 (13)	11.1 (41)	2,281	4.2 (18)	8.6 (57)	11.1 (8)	903	13.7 (6)	11.7 (6)	1,576	14.0 (40)
Tehama	417	5.8 (12)	9.1 (19)	1,892	3.3 (3)	6.7 (39)	13.4 (24)	852	15.5 (20)	15.0 (24)	1,470	15.4 (52)
Trinity	49	* *	* *	372	5.1 (39)	8.9 (60)	14.5 (37)	176	22.2 (64)	15.9 (34)	211	10.4 (12)
Tulare	2,725	6.3 (19)	10.3 (31)	17,196	5.4 (43)	4.7 (9)	16.9 (56)	6,531	16.5 (37)	16.1 (40)	11,845	12.2 (28)
Tuolumne	55	* *	* *	749	5.7 (50)	6.5 (34)	11.6 (10)	302	12.6 (3)	12.3 (9)	398	9.8 (6)
Ventura	3,441	5.9 (13)	10.6 (35)	14,421	3.6 (6)	5.0 (16)	14.3 (35)	5,377	16.0 (32)	15.9 (34)	10,384	11.6 (20)
Yolo	413	4.6 (4)	13.6 (54)	3,597	3.8 (7)	9.2 (63)	11.7 (13)	1,386	15.2 (14)	14.3 (17)	1,976	16.5 (59)
Yuba	313	7.3 (34)	11.5 (44)	1,526	5.2 (41)	7.0 (43)	11.6 (10)	575	13.2 (4)	12.0 (8)	977	14.4 (45)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**1999 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
<b>California</b>	<b>162,300</b>	<b>6.9</b>	<b>9.6</b>	<b>897,904</b>	<b>5.0</b>	<b>5.2</b>	<b>15.6</b>	<b>375,498</b>	<b>16.3</b>	<b>15.9</b>	<b>582,213</b>	<b>13.0</b>
<b>Nation (Prior Year)</b>	<b>1,307,687</b>	<b>8.9</b>	<b>8.5</b>	<b>5,365,056</b>	<b>6.0</b>	<b>5.9</b>	<b>12.0</b>	<b>2,320,924</b>	<b>14.9</b>	<b>12.2</b>	<b>3,864,299</b>	<b>14.5</b>

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**1999 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged 5 to <20 Years** (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alameda	16,589	7.6 (48)	3.2 (54)	16.8 (20)	18.4 (24)	10,259	15.3 (57)
Alpine	233	10.7 (64)	1.3 (4)	16.7 (19)	20.6 (51)	184	11.4 (32)
Amador	674	4.9 (5)	2.7 (33)	15.4 (5)	18.1 (20)	521	10.4 (20)
Butte	3,512	7.7 (50)	3.4 (56)	15.5 (6)	15.9 (11)	3,465	10.3 (16)
Calaveras	726	8.5 (55)	2.3 (20)	18.7 (53)	18.0 (19)	490	11.2 (29)
City Of Be	319	7.2 (38)	2.5 (28)	17.6 (32)	21.9 (61)	232	22.4 (63)
Colusa	807	6.8 (29)	1.6 (7)	19.2 (58)	21.8 (60)	701	8.7 (4)
Contra Cos	6,026	7.4 (43)	2.3 (20)	18.2 (45)	19.8 (45)	4,315	15.3 (57)
Del Norte	501	5.8 (10)	2.8 (38)	21.8 (64)	15.2 (8)	435	17.9 (61)
El Dorado	2,293	5.8 (10)	2.8 (38)	16.6 (16)	18.4 (24)	1,893	10.0 (11)
Fresno	29,285	8.6 (56)	2.7 (33)	16.8 (20)	19.3 (37)	22,989	10.4 (20)
Glenn	621	7.4 (43)	2.6 (29)	18.5 (51)	20.0 (48)	631	12.4 (35)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**1999 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged 5 to <20 Years** (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Humboldt	1,841	6.0 (15)	2.7 (33)	18.4 (50)	18.6 (28)	1,397	12.9 (41)
Imperial	3,124	6.4 (20)	2.3 (20)	19.0 (56)	24.8 (64)	2,932	5.5 (1)
Inyo	708	6.6 (26)	1.7 (10)	18.6 (52)	21.9 (61)	524	18.5 (62)
Kern	12,213	6.9 (32)	2.9 (44)	16.6 (16)	20.3 (49)	9,195	11.0 (27)
Kings	1,829	5.4 (8)	3.0 (51)	16.5 (14)	19.8 (45)	1,845	15.0 (56)
L.A. East	39,873	6.5 (24)	2.6 (29)	18.2 (45)	19.3 (37)	38,347	11.6 (33)
L.A. North	48,203	6.8 (29)	2.3 (20)	18.3 (48)	18.2 (22)	48,753	9.2 (7)
L.A. Other	48,126	4.9 (5)	2.8 (38)	17.7 (35)	21.7 (57)	36,215	13.0 (43)
L.A. South	65,945	6.5 (24)	2.9 (44)	18.1 (43)	18.7 (31)	64,631	10.6 (23)
L.A. West	18,852	5.9 (12)	2.8 (38)	18.0 (40)	18.6 (28)	16,690	13.2 (45)
Lake	777	7.6 (48)	1.9 (14)	15.6 (7)	19.9 (47)	646	12.8 (39)
Lassen	410	4.6 (3)	1.2 (3)	15.1 (4)	13.7 (3)	311	11.3 (30)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**1999 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged 5 to <20 Years** (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Long Beach	8,797	5.9 (12)	2.9 (44)	17.4 (28)	19.4 (39)	7,052	13.2 (45)
Madera	2,761	6.6 (26)	2.9 (44)	16.6 (16)	19.1 (35)	2,827	10.0 (11)
Marin	1,018	7.2 (38)	1.1 (1)	17.7 (35)	20.7 (52)	619	12.4 (35)
Mariposa	185	4.9 (5)	1.6 (7)	16.8 (20)	11.4 (1)	175	8.6 (3)
Mendocino	1,398	6.4 (20)	1.4 (6)	18.9 (54)	20.5 (50)	859	13.2 (45)
Merced	5,523	10.3 (62)	2.2 (18)	17.5 (29)	19.4 (39)	5,469	13.7 (50)
Modoc	212	10.4 (63)	2.4 (24)	16.5 (14)	21.7 (57)	170	8.8 (5)
Mono	114	8.8 (58)	2.6 (29)	10.5 (1)	18.4 (24)	84	* *
Monterey	5,887	7.2 (38)	1.6 (7)	19.2 (58)	22.6 (63)	6,261	9.7 (8)
Napa	1,030	7.9 (51)	2.4 (24)	17.9 (39)	19.0 (33)	557	13.1 (44)
Nevada	545	8.8 (58)	4.6 (62)	12.8 (2)	14.3 (4)	470	8.1 (2)
Orange	61,945	7.1 (37)	2.9 (44)	17.6 (32)	17.9 (17)	54,910	14.0 (52)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**1999 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged 5 to <20 Years** (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Pasadena	2,344	5.7 (9)	2.9 (44)	17.1 (25)	17.7 (15)	2,127	10.3 (16)
Placer	1,098	8.2 (53)	3.8 (60)	17.6 (32)	16.5 (12)	756	10.6 (23)
Plumas	464	6.3 (19)	4.7 (63)	19.2 (58)	19.6 (44)	300	10.0 (11)
Riverside	24,446	6.2 (17)	2.8 (38)	17.2 (26)	18.4 (24)	21,676	12.6 (37)
Sacramento	21,093	8.3 (54)	3.5 (57)	15.6 (7)	15.6 (9)	14,400	16.2 (60)
San Benito	1,197	7.2 (38)	2.2 (18)	18.9 (54)	21.7 (57)	1,066	8.9 (6)
San Bernar	38,817	6.4 (20)	3.2 (54)	16.9 (23)	18.1 (20)	30,308	12.9 (41)
San Diego	34,093	7.0 (36)	2.4 (24)	17.7 (35)	19.4 (39)	29,750	10.8 (26)
San Franci	6,629	8.1 (52)	3.7 (59)	16.3 (12)	16.7 (13)	3,190	11.1 (28)
San Joaqui	9,916	8.9 (60)	3.1 (52)	16.3 (12)	17.9 (17)	6,667	12.8 (39)
San Luis O	2,041	7.3 (42)	2.4 (24)	17.8 (38)	18.2 (22)	1,125	10.1 (14)
San Mateo	5,457	9.2 (61)	1.8 (11)	19.3 (62)	21.3 (55)	3,772	9.8 (9)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**1999 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged 5 to <20 Years** (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Santa Barb	4,900	7.5 (47)	2.1 (16)	19.3 (62)	20.9 (53)	4,807	10.7 (25)
Santa Clar	11,920	8.6 (56)	3.6 (58)	16.2 (11)	18.6 (28)	7,432	13.6 (49)
Santa Cruz	3,947	6.2 (17)	1.8 (11)	18.1 (43)	21.5 (56)	3,149	10.3 (16)
Shasta	2,052	5.9 (12)	3.1 (52)	17.3 (27)	15.0 (7)	1,410	10.1 (14)
Sierra	63	* *	* *	* *	* *	40	* *
Siskiyou	842	3.9 (1)	1.3 (4)	15.6 (7)	14.3 (4)	717	11.3 (30)
Solano	3,276	6.1 (16)	2.9 (44)	17.0 (24)	19.5 (43)	2,501	15.8 (59)
Sonoma	2,744	6.8 (29)	1.9 (14)	19.1 (57)	19.4 (39)	1,658	13.5 (48)
Stanislaus	7,806	6.9 (32)	2.7 (33)	18.0 (40)	18.8 (32)	8,040	14.6 (55)
Sutter	1,045	6.4 (20)	4.5 (61)	15.9 (10)	14.7 (6)	964	14.0 (52)
Tehama	992	7.4 (43)	1.8 (11)	18.0 (40)	17.5 (14)	916	12.3 (34)
Trinity	177	4.0 (2)	1.1 (1)	17.5 (29)	15.8 (10)	132	9.8 (9)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**1999 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged 5 to <20 Years** (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Tulare	9,457	7.4 (43)	2.8 (38)	17.5 (29)	19.0 (33)	9,049	10.3 (16)
Tuolumne	364	6.9 (32)	4.9 (64)	14.0 (3)	13.5 (2)	212	12.7 (38)
Ventura	7,175	4.7 (4)	2.6 (29)	18.3 (48)	21.2 (54)	7,161	10.5 (22)
Yolo	1,951	6.6 (26)	2.1 (16)	18.2 (45)	19.1 (35)	1,416	13.7 (50)
Yuba	897	6.9 (32)	2.7 (33)	19.2 (58)	17.7 (15)	812	14.3 (54)
<b>California</b>	<b>600,105</b>	<b>6.8</b>	<b>2.8</b>	<b>17.5</b>	<b>18.9</b>	<b>512,607</b>	<b>11.9</b>

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.