

**Table 6B**  
**2000 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alameda	4,310	7.3 (34)	9.7 (25)	23,680	6.1 (61)	5.8 (30)	15.7 (43)	8,977	16.4 (33)	16.8 (43)	10,375	14.1 (39)
Alpine	23	* *	* *	199	7.0 (64)	4.0 (3)	19.1 (62)	99	* *	* *	122	16.4 (52)
Amador	141	9.9 (54)	11.3 (41)	775	4.1 (18)	7.1 (43)	13.9 (30)	322	13.7 (3)	14.0 (20)	413	12.3 (27)
Butte	1,313	7.8 (43)	8.5 (9)	6,087	4.2 (19)	7.0 (40)	11.5 (19)	2,524	14.7 (11)	12.1 (7)	4,431	10.3 (9)
Calaveras	203	6.4 (19)	9.9 (27)	1,067	6.2 (62)	5.5 (26)	15.1 (37)	403	16.9 (44)	12.4 (11)	571	12.6 (31)
City Of Be	72	* *	* *	385	6.0 (59)	7.0 (40)	16.4 (51)	148	16.9 (44)	17.6 (50)	177	24.3 (63)
Colusa	296	4.7 (4)	9.8 (26)	1,905	4.4 (24)	8.7 (57)	16.3 (50)	790	15.6 (20)	19.2 (57)	1,103	17.3 (58)
Contra Cos	1,871	7.0 (28)	10.0 (28)	11,723	5.1 (41)	6.2 (32)	15.1 (37)	4,348	19.3 (62)	18.2 (55)	5,090	13.8 (37)
Del Norte	179	6.7 (23)	13.4 (51)	904	5.1 (41)	5.0 (18)	16.2 (47)	375	18.7 (60)	16.5 (38)	627	17.7 (60)
El Dorado	452	8.0 (44)	8.0 (7)	2,331	4.7 (28)	8.5 (54)	11.3 (16)	896	15.8 (23)	12.1 (7)	1,339	11.0 (15)
Fresno	7,339	7.2 (33)	8.7 (12)	39,059	5.8 (56)	4.0 (3)	16.7 (54)	16,507	15.9 (26)	16.2 (33)	27,010	12.1 (25)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Glenn	281	7.5 (37)	10.7 (37)	1,207	3.1 (4)	5.1 (20)	11.1 (13)	516	17.1 (49)	13.2 (16)	942	12.5 (30)
Humboldt	520	6.9 (26)	14.8 (55)	3,602	4.7 (28)	5.4 (23)	13.5 (26)	1,462	17.0 (48)	16.4 (35)	2,114	16.7 (56)
Imperial	962	5.5 (10)	12.0 (46)	5,790	2.7 (2)	8.3 (53)	15.5 (42)	2,251	15.9 (26)	17.2 (45)	3,947	6.3 (1)
Inyo	103	1.9 (1)	14.6 (54)	742	4.6 (27)	4.9 (16)	19.5 (63)	314	15.9 (26)	20.4 (60)	438	16.0 (50)
Kern	5,733	7.1 (31)	10.6 (36)	26,214	4.9 (36)	5.1 (20)	16.0 (46)	10,835	16.3 (31)	15.8 (30)	15,864	8.7 (5)
Kings	547	6.8 (24)	12.4 (47)	2,866	5.0 (39)	3.6 (1)	17.6 (58)	1,067	16.9 (44)	15.1 (28)	1,644	18.7 (61)
L.A. East	16,679	6.1 (14)	9.1 (16)	59,867	5.3 (45)	4.4 (9)	17.8 (60)	22,155	15.8 (23)	17.5 (47)	36,343	15.2 (47)
L.A. North	11,123	6.3 (16)	9.5 (22)	54,818	5.0 (39)	4.8 (15)	16.5 (53)	22,023	16.5 (37)	17.4 (46)	33,382	12.1 (25)
L.A. Other	7,312	8.0 (44)	9.1 (16)	77,807	4.3 (20)	4.0 (3)	18.2 (61)	37,585	15.8 (23)	17.6 (50)	52,261	13.1 (34)
L.A. South	17,389	7.0 (28)	9.2 (18)	75,194	5.9 (57)	4.0 (3)	20.0 (64)	31,499	17.1 (49)	19.3 (58)	50,038	12.8 (32)
L.A. West	5,579	7.0 (28)	9.6 (24)	23,420	5.4 (48)	4.3 (7)	17.7 (59)	8,799	16.6 (39)	18.0 (54)	13,453	14.2 (42)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Lake	308	7.5 (37)	13.0 (49)	1,308	4.3 (20)	7.4 (46)	11.3 (16)	512	14.8 (14)	12.5 (12)	860	11.3 (18)
Lassen	169	7.7 (39)	9.5 (22)	590	3.9 (15)	9.8 (61)	6.9 (1)	220	14.1 (5)	5.5 (1)	296	10.8 (14)
Long Beach	2,902	6.6 (21)	8.9 (13)	13,420	5.3 (45)	4.3 (7)	16.9 (56)	5,394	16.9 (44)	17.5 (47)	7,448	17.4 (59)
Madera	1,410	7.7 (39)	8.6 (10)	5,120	5.7 (51)	4.6 (11)	15.3 (39)	2,170	15.9 (26)	14.5 (22)	4,220	13.5 (36)
Marin	107	5.6 (11)	13.1 (50)	2,025	2.8 (3)	6.6 (37)	10.8 (11)	706	14.4 (10)	16.4 (35)	1,101	8.4 (3)
Mariposa	42	* *	* *	217	2.3 (1)	6.5 (35)	7.8 (4)	80	* *	* *	157	11.5 (20)
Mendocino	622	5.1 (6)	10.1 (30)	2,377	3.5 (7)	9.3 (60)	10.6 (10)	898	17.7 (56)	14.6 (23)	993	17.0 (57)
Merced	2,188	8.4 (50)	10.1 (30)	9,728	5.7 (51)	3.9 (2)	15.7 (43)	3,758	18.5 (59)	17.1 (44)	6,929	15.1 (45)
Modoc	62	* *	* *	262	3.8 (12)	9.9 (62)	9.5 (8)	115	17.4 (54)	12.2 (9)	170	11.8 (23)
Mono	88	* *	* *	310	4.8 (35)	13.2 (64)	7.7 (3)	116	12.9 (1)	10.3 (4)	167	26.3 (64)
Monterey	942	5.3 (7)	10.0 (28)	8,278	4.3 (20)	4.9 (16)	15.9 (45)	3,240	19.0 (61)	20.5 (61)	6,266	12.3 (27)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Napa	190	4.7 (4)	11.6 (44)	1,832	3.8 (12)	8.1 (49)	11.2 (14)	559	17.9 (57)	14.7 (25)	664	16.1 (51)
Nevada	266	4.1 (2)	11.3 (41)	1,063	3.6 (10)	11.7 (63)	7.2 (2)	341	14.7 (11)	8.8 (2)	530	11.5 (20)
Orange	10,880	6.3 (16)	9.2 (18)	75,651	4.5 (26)	5.4 (23)	16.2 (47)	32,710	16.1 (30)	16.7 (41)	50,181	14.6 (44)
Pasadena	891	8.5 (52)	3.1 (1)	3,577	4.9 (36)	6.4 (33)	14.2 (32)	1,366	16.6 (39)	13.4 (18)	2,162	8.6 (4)
Placer	397	4.3 (3)	11.6 (44)	1,924	6.0 (59)	7.6 (48)	10.8 (11)	638	14.3 (8)	16.6 (40)	1,157	11.6 (22)
Plumas	89	* *	* *	456	4.4 (24)	6.8 (38)	8.3 (5)	184	14.1 (5)	10.3 (4)	227	8.8 (6)
Riverside	7,710	6.9 (26)	7.9 (6)	42,605	5.4 (48)	4.7 (12)	16.2 (47)	17,919	15.7 (22)	15.9 (32)	27,049	14.1 (39)
Sacramento	3,009	8.3 (48)	9.4 (21)	27,733	5.2 (44)	7.0 (40)	13.1 (25)	11,357	15.3 (17)	14.6 (23)	16,472	15.2 (47)
San Benito	175	7.4 (36)	10.3 (33)	1,497	3.8 (12)	5.4 (23)	16.7 (54)	621	15.5 (19)	20.8 (62)	1,052	11.4 (19)
San Bernar	9,183	6.8 (24)	8.2 (8)	56,569	5.9 (57)	5.5 (26)	15.4 (41)	25,212	14.9 (15)	14.4 (21)	33,525	16.4 (52)
San Diego	9,613	6.5 (20)	11.0 (40)	58,439	4.7 (28)	6.5 (35)	13.7 (29)	23,599	15.4 (18)	15.0 (27)	38,197	11.0 (15)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
San Franci	736	8.3 (48)	7.5 (2)	8,343	5.3 (45)	5.0 (18)	14.7 (36)	3,148	17.2 (53)	17.7 (53)	1,929	10.6 (12)
San Joaqui	4,044	8.0 (44)	9.3 (20)	18,244	6.7 (63)	5.7 (29)	16.4 (51)	7,303	17.1 (49)	18.6 (56)	10,462	10.4 (11)
San Luis O	561	7.7 (39)	10.7 (37)	4,066	3.9 (15)	8.1 (49)	10.4 (9)	1,556	16.3 (31)	13.2 (16)	1,967	10.7 (13)
San Mateo	999	6.6 (21)	7.7 (4)	8,354	4.7 (28)	7.3 (45)	14.4 (33)	3,044	17.1 (49)	19.8 (59)	4,033	7.8 (2)
Santa Barb	2,739	10.4 (55)	7.5 (2)	11,158	5.5 (50)	6.0 (31)	14.5 (35)	4,130	16.8 (41)	17.5 (47)	7,719	10.2 (8)
Santa Clar	2,244	5.7 (12)	9.0 (15)	22,191	4.7 (28)	6.9 (39)	13.6 (27)	7,859	15.6 (20)	16.7 (41)	10,789	15.1 (45)
Santa Cruz	1,356	6.3 (16)	10.9 (39)	10,665	4.0 (17)	4.7 (12)	15.3 (39)	3,886	18.4 (58)	17.6 (50)	6,161	12.4 (29)
Shasta	802	8.4 (50)	10.3 (33)	3,321	5.7 (51)	5.6 (28)	14.1 (31)	1,393	16.8 (41)	12.3 (10)	2,123	11.0 (15)
Sierra	14	* *	* *	57	* *	* *	* *	25	* *	* *	24	* *
Siskiyou	217	10.6 (56)	7.8 (5)	1,106	4.7 (28)	7.4 (46)	11.8 (22)	463	13.4 (2)	13.4 (18)	709	16.5 (54)
Solano	1,407	8.1 (47)	8.6 (10)	6,860	4.9 (36)	8.1 (49)	12.7 (24)	2,424	16.4 (33)	15.8 (30)	3,411	14.3 (43)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Sonoma	1,181	5.8 (13)	12.5 (48)	5,756	3.2 (5)	8.7 (57)	11.3 (16)	1,891	15.0 (16)	15.5 (29)	2,394	13.3 (35)
Stanislaus	3,008	9.6 (53)	8.9 (13)	13,872	4.3 (20)	8.5 (54)	13.6 (27)	5,435	16.4 (33)	16.5 (38)	10,254	16.6 (55)
Sutter	533	6.2 (15)	16.7 (56)	2,252	3.5 (7)	8.5 (54)	11.2 (14)	859	14.0 (4)	12.9 (14)	1,404	15.4 (49)
Tehama	436	7.1 (31)	10.3 (33)	1,880	3.4 (6)	8.1 (49)	11.5 (19)	752	17.4 (54)	12.8 (13)	1,329	19.6 (62)
Trinity	17	* *	* *	228	5.7 (51)	4.4 (9)	9.2 (7)	113	14.2 (7)	9.7 (3)	125	12.8 (32)
Tulare	2,938	7.3 (34)	10.1 (30)	17,212	5.7 (51)	4.7 (12)	17.4 (57)	6,438	16.5 (37)	16.4 (35)	11,460	14.1 (39)
Tuolumne	40	* *	* *	784	4.7 (28)	6.4 (33)	8.5 (6)	306	14.7 (11)	13.1 (15)	403	9.9 (7)
Ventura	3,602	5.3 (7)	11.4 (43)	14,623	3.7 (11)	5.1 (20)	14.4 (33)	5,518	16.8 (41)	16.3 (34)	9,870	10.3 (9)
Yolo	790	5.3 (7)	13.4 (51)	3,443	3.5 (7)	9.0 (59)	12.2 (23)	1,247	16.4 (33)	14.9 (26)	1,817	13.8 (37)
Yuba	350	7.7 (39)	13.4 (51)	1,562	5.1 (41)	7.2 (44)	11.5 (19)	565	14.3 (8)	12.0 (6)	877	12.0 (24)

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

<b>Comparison of Growth and Anemia Indicators by County</b> (3,4)												
<b>County</b>	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
<b>California</b>	<b>161,684</b>	<b>6.9</b>	<b>9.4</b>	<b>890,600</b>	<b>5.1</b>	<b>5.3</b>	<b>16.1</b>	<b>363,965</b>	<b>16.2</b>	<b>16.7</b>	<b>550,767</b>	<b>13.3</b>
<b>Nation (Prior Year)</b>	<b>1,301,847</b>	<b>8.9</b>	<b>8.4</b>	<b>5,149,796</b>	<b>5.9</b>	<b>5.7</b>	<b>12.3</b>	<b>2,187,151</b>	<b>15.0</b>	<b>12.7</b>	<b>3,645,746</b>	<b>14.5</b>

- (1) Reporting period is January 1 through December 31.  
(2) Analyses based on one record per child.  
(3) Excludes records with unknown data or errors.  
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.  
(5) Infants born during the reporting period included in the analysis.  
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.  
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.  
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.  
\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged 5 to <20 Years** (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alameda	16,688	7.8 (47)	3.1 (49)	17.6 (32)	19.3 (30)	9,968	15.8 (53)
Alpine	106	11.3 (63)	1.9 (9)	18.9 (47)	22.6 (60)	96	* *
Amador	448	6.3 (19)	3.3 (58)	15.8 (8)	18.1 (16)	366	11.5 (27)
Butte	3,204	6.9 (29)	3.0 (45)	16.1 (9)	17.3 (11)	3,145	10.2 (13)
Calaveras	831	8.5 (53)	2.3 (22)	15.4 (5)	19.3 (30)	677	12.4 (36)
City Of Be	198	7.6 (41)	3.0 (45)	23.7 (63)	19.2 (25)	132	22.0 (62)
Colusa	1,273	6.8 (27)	1.6 (7)	17.6 (32)	20.8 (51)	949	13.0 (41)
Contra Cos	6,789	6.6 (23)	2.4 (25)	18.2 (42)	20.8 (51)	4,505	15.3 (50)
Del Norte	298	7.0 (32)	2.0 (13)	20.5 (61)	17.8 (15)	278	14.0 (46)
El Dorado	1,240	7.7 (46)	3.5 (60)	15.3 (4)	18.5 (19)	1,113	10.7 (16)
Fresno	29,795	9.2 (58)	2.6 (30)	17.1 (25)	20.1 (43)	23,588	11.1 (21)
Glenn	465	8.0 (50)	2.6 (30)	20.2 (56)	18.7 (20)	463	11.4 (26)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged 5 to <20 Years** (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Humboldt	1,603	6.0 (12)	1.2 (2)	19.0 (49)	20.0 (41)	1,152	16.3 (56)
Imperial	3,155	5.8 (7)	2.6 (30)	17.5 (31)	22.9 (61)	2,972	5.4 (2)
Inyo	409	7.1 (33)	3.4 (59)	17.8 (36)	21.8 (57)	300	19.3 (60)
Kern	12,961	6.5 (22)	3.1 (49)	16.6 (14)	20.2 (45)	9,787	10.8 (17)
Kings	1,546	5.8 (7)	3.0 (45)	15.7 (6)	20.4 (47)	1,522	15.8 (53)
L.A. East	37,548	6.6 (23)	2.7 (37)	18.1 (39)	20.2 (45)	36,222	12.9 (40)
L.A. North	44,293	6.7 (26)	2.6 (30)	18.0 (38)	18.8 (22)	44,206	9.9 (10)
L.A. Other	48,906	4.9 (4)	2.7 (37)	17.6 (32)	22.4 (58)	36,350	12.5 (37)
L.A. South	65,856	7.3 (36)	2.6 (30)	19.1 (52)	19.5 (33)	64,196	10.8 (17)
L.A. West	16,068	6.1 (15)	2.6 (30)	19.1 (52)	19.6 (36)	14,007	14.2 (47)
Lake	651	7.5 (40)	3.5 (60)	16.4 (11)	19.5 (33)	579	8.3 (5)
Lassen	341	7.6 (41)	2.1 (17)	16.7 (15)	17.0 (7)	246	8.9 (6)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged 5 to <20 Years** (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Long Beach	8,234	6.0 (12)	2.7 (37)	17.3 (28)	19.9 (40)	6,436	13.8 (45)
Madera	3,102	7.3 (36)	2.6 (30)	16.9 (19)	19.2 (25)	3,173	11.1 (21)
Marin	808	9.3 (59)	1.9 (9)	20.2 (56)	21.2 (55)	529	9.1 (8)
Mariposa	170	8.2 (51)	2.4 (25)	12.9 (1)	12.9 (3)	156	10.9 (19)
Mendocino	1,296	7.1 (33)	1.8 (8)	16.9 (19)	21.0 (54)	840	14.6 (49)
Merced	5,097	8.8 (55)	2.3 (22)	17.0 (23)	20.1 (43)	5,205	17.0 (59)
Modoc	183	8.7 (54)	0.5 (1)	20.8 (62)	19.1 (24)	156	4.5 (1)
Mono	86	* *	* *	* *	* *	88	* *
Monterey	3,635	7.9 (48)	1.5 (5)	20.2 (56)	22.4 (58)	3,889	11.6 (28)
Napa	882	6.1 (15)	2.2 (19)	18.9 (47)	20.0 (41)	493	19.3 (60)
Nevada	426	5.9 (10)	1.4 (3)	13.6 (2)	17.1 (8)	323	9.9 (10)
Orange	52,464	7.3 (36)	3.1 (49)	18.1 (39)	19.2 (25)	44,571	13.7 (44)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged 5 to <20 Years** (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Pasadena	1,639	4.5 (3)	3.1 (49)	17.0 (23)	17.3 (11)	1,503	11.6 (28)
Placer	979	10.0 (61)	4.0 (63)	14.2 (3)	19.6 (36)	794	11.3 (24)
Plumas	277	7.6 (41)	1.4 (3)	17.3 (28)	16.2 (5)	203	5.9 (3)
Riverside	28,933	6.4 (20)	3.0 (45)	16.7 (15)	19.2 (25)	25,438	15.9 (55)
Sacramento	20,436	7.9 (48)	3.1 (49)	15.7 (6)	17.2 (9)	13,931	16.6 (57)
San Benito	918	7.6 (41)	2.0 (13)	20.2 (56)	23.3 (63)	878	9.1 (8)
San Bernar	41,237	6.4 (20)	3.2 (57)	16.9 (19)	19.0 (23)	31,923	14.2 (47)
San Diego	30,324	6.6 (23)	2.7 (37)	17.9 (37)	19.5 (33)	27,648	13.1 (42)
San Franci	6,308	8.8 (55)	3.1 (49)	16.2 (10)	18.3 (18)	2,714	10.9 (19)
San Joaqui	9,789	9.4 (60)	3.1 (49)	16.4 (11)	18.7 (20)	6,340	12.2 (33)
San Luis O	1,781	7.1 (33)	2.2 (19)	19.0 (49)	18.2 (17)	1,002	11.1 (21)
San Mateo	5,112	9.1 (57)	2.0 (13)	20.3 (60)	21.3 (56)	3,601	9.0 (7)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged 5 to <20 Years** (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Santa Barb	4,201	7.6 (41)	2.0 (13)	19.5 (55)	20.9 (53)	4,196	10.5 (15)
Santa Clar	11,209	8.2 (51)	2.9 (44)	16.8 (17)	19.2 (25)	7,292	13.1 (42)
Santa Cruz	4,682	5.9 (10)	1.5 (5)	19.1 (52)	23.2 (62)	3,929	11.9 (30)
Shasta	1,600	6.2 (18)	2.4 (25)	18.1 (39)	16.1 (4)	1,237	10.4 (14)
Sierra	44	* *	* *	* *	* *	34	* *
Siskiyou	712	4.1 (1)	2.7 (37)	18.3 (43)	17.7 (14)	566	12.0 (32)
Solano	3,292	6.1 (15)	3.1 (49)	16.5 (13)	20.6 (49)	2,474	16.9 (58)
Sonoma	2,597	6.9 (29)	2.4 (25)	18.4 (44)	17.2 (9)	1,535	12.6 (38)
Stanislaus	7,396	6.9 (29)	2.4 (25)	18.4 (44)	19.7 (39)	7,579	15.6 (52)
Sutter	936	5.7 (6)	3.6 (62)	19.0 (49)	16.9 (6)	823	11.3 (24)
Tehama	935	6.8 (27)	2.1 (17)	17.1 (25)	19.3 (30)	858	15.4 (51)
Trinity	160	5.0 (5)	1.9 (9)	18.8 (46)	10.6 (2)	121	6.6 (4)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2000 Pediatric Nutrition Surveillance** (1)  
**California**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged 5 to <20 Years** (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Tulare	9,285	7.3 (36)	2.3 (22)	17.2 (27)	19.6 (36)	8,616	12.2 (33)
Tuolumne	364	5.8 (7)	1.9 (9)	16.8 (17)	10.2 (1)	201	11.9 (30)
Ventura	6,040	4.4 (2)	2.7 (37)	17.6 (32)	20.7 (50)	5,855	10.0 (12)
Yolo	1,749	6.0 (12)	2.2 (19)	16.9 (19)	20.4 (47)	1,324	12.2 (33)
Yuba	830	10.1 (62)	2.7 (37)	17.3 (28)	17.6 (13)	772	12.6 (38)
<b>California</b>	<b>574,820</b>	<b>6.9</b>	<b>2.8</b>	<b>17.7</b>	<b>19.7</b>	<b>486,065</b>	<b>12.6</b>

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.