

Table 6B
2001 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alameda	3,203	7.4 (41)	10.2 (31)	18,499	6.2 (58)	6.0 (28)	15.3 (44)	6,235	16.5 (34)	16.4 (33)	7,313	12.7 (26)
Alpine	25	* *	* *	147	7.5 (63)	8.2 (47)	17.0 (55)	59	* *	* *	84	* *
Amador	136	14.0 (53)	9.6 (25)	588	4.4 (20)	9.7 (57)	13.4 (27)	231	14.7 (9)	14.3 (14)	260	9.2 (6)
Butte	1,188	6.1 (20)	10.3 (33)	5,852	5.1 (38)	7.2 (39)	11.5 (14)	2,404	14.1 (5)	12.0 (6)	3,954	9.8 (11)
Calaveras	134	5.2 (8)	14.9 (54)	806	4.6 (24)	7.3 (40)	15.0 (41)	322	13.0 (2)	16.1 (29)	457	12.5 (22)
City Of Be	41	* *	* *	258	5.0 (35)	4.7 (14)	15.9 (48)	98	* *	* *	119	16.0 (50)
Colusa	224	4.0 (3)	8.0 (8)	1,435	3.9 (10)	9.2 (52)	14.1 (34)	541	17.7 (48)	17.0 (40)	704	13.1 (30)
Contra Cos	1,182	7.4 (41)	10.8 (35)	8,865	5.8 (48)	6.3 (29)	14.5 (38)	2,873	17.5 (46)	16.4 (33)	3,192	14.2 (38)
Del Norte	149	3.4 (2)	12.8 (50)	829	5.9 (51)	4.2 (7)	16.6 (52)	346	20.5 (57)	14.7 (18)	607	13.7 (33)
El Dorado	249	10.4 (50)	14.1 (51)	1,966	4.9 (30)	9.4 (54)	9.7 (5)	700	13.9 (4)	11.9 (5)	991	12.6 (25)
Fresno	6,235	7.0 (34)	9.1 (20)	34,408	5.6 (45)	4.2 (7)	15.5 (46)	13,903	15.9 (18)	15.6 (25)	21,819	12.9 (29)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Glenn	257	4.3 (5)	10.9 (38)	1,119	3.0 (2)	5.9 (26)	13.9 (31)	459	16.3 (29)	16.8 (38)	806	9.7 (10)
Humboldt	583	4.1 (4)	14.4 (53)	3,382	4.8 (28)	4.8 (16)	13.7 (29)	1,387	19.0 (54)	16.2 (31)	2,031	16.4 (53)
Imperial	913	6.4 (23)	10.8 (35)	5,563	3.0 (2)	8.1 (46)	14.0 (32)	2,207	15.0 (11)	15.6 (25)	3,445	9.3 (8)
Inyo	76	* *	* *	485	6.4 (59)	4.3 (10)	15.5 (46)	189	15.3 (14)	18.5 (52)	208	10.6 (13)
Kern	4,182	5.2 (8)	11.8 (45)	23,238	4.9 (30)	5.2 (21)	16.9 (53)	9,340	16.4 (31)	16.6 (37)	13,311	10.6 (13)
Kings	381	6.8 (29)	9.4 (23)	2,314	5.0 (35)	3.9 (1)	18.8 (63)	863	17.7 (48)	17.1 (42)	1,381	15.6 (49)
L.A. East	15,049	6.0 (18)	8.8 (16)	53,292	5.6 (45)	4.4 (11)	18.0 (61)	19,448	16.2 (27)	17.5 (46)	28,386	16.3 (52)
L.A. North	9,655	5.9 (15)	8.4 (11)	39,396	5.5 (43)	4.7 (14)	17.1 (56)	15,492	16.7 (37)	18.1 (50)	21,851	12.3 (21)
L.A. Other	4,914	8.0 (45)	8.5 (13)	86,766	3.9 (10)	4.0 (2)	18.0 (61)	40,315	16.3 (29)	17.7 (48)	55,415	13.6 (32)
L.A. South	14,895	7.1 (37)	8.9 (19)	58,129	6.1 (57)	4.0 (2)	19.3 (64)	22,436	17.1 (41)	19.0 (55)	32,882	12.8 (27)
L.A. West	4,308	6.5 (26)	9.2 (21)	18,613	5.5 (43)	4.2 (7)	17.3 (57)	6,955	16.6 (36)	17.3 (43)	9,662	14.2 (38)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Lake	208	6.7 (27)	12.5 (48)	1,064	4.5 (21)	6.3 (29)	13.2 (24)	435	20.5 (57)	16.1 (29)	734	14.0 (35)
Lassen	127	15.0 (54)	8.7 (15)	435	3.9 (10)	11.7 (61)	9.4 (4)	134	16.4 (31)	9.0 (1)	190	8.9 (5)
Long Beach	2,134	6.4 (23)	7.3 (4)	11,434	4.9 (30)	4.4 (11)	17.3 (57)	4,441	17.2 (42)	17.6 (47)	5,368	17.3 (56)
Madera	1,104	6.9 (32)	7.0 (3)	4,752	4.7 (27)	4.8 (16)	13.3 (25)	1,954	16.0 (21)	11.7 (4)	3,649	15.3 (48)
Marin	75	* *	* *	1,733	3.4 (5)	7.5 (41)	9.9 (7)	542	14.4 (8)	15.7 (28)	953	6.7 (1)
Mariposa	37	* *	* *	186	5.4 (42)	11.8 (62)	5.9 (1)	70	* *	* *	118	11.0 (16)
Mendocino	553	5.4 (11)	10.1 (29)	2,043	3.5 (6)	10.4 (59)	11.6 (16)	684	18.9 (52)	15.1 (20)	782	16.4 (53)
Merced	1,887	6.0 (18)	10.0 (28)	9,191	5.3 (41)	4.1 (4)	15.1 (42)	3,478	17.3 (44)	17.7 (48)	6,392	17.3 (56)
Modoc	27	* *	* *	175	5.7 (47)	16.0 (64)	10.9 (12)	85	* *	* *	116	8.6 (4)
Mono	65	* *	* *	278	5.8 (48)	13.7 (63)	6.5 (2)	80	* *	* *	141	20.6 (61)
Monterey	879	6.9 (32)	10.8 (35)	6,475	4.5 (21)	4.6 (13)	14.3 (36)	2,532	18.2 (50)	18.7 (54)	4,364	11.3 (19)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Napa	186	7.0 (34)	10.2 (31)	1,811	3.5 (6)	8.5 (49)	10.0 (8)	570	17.2 (42)	14.7 (18)	622	19.9 (59)
Nevada	233	8.6 (47)	11.6 (43)	944	4.8 (28)	9.9 (58)	7.1 (3)	310	16.8 (40)	11.3 (3)	439	10.3 (12)
Orange	8,293	6.8 (29)	8.0 (8)	67,474	4.0 (16)	5.4 (22)	15.9 (48)	27,778	16.1 (24)	16.4 (33)	41,556	13.9 (34)
Pasadena	773	6.2 (22)	6.5 (1)	2,701	4.6 (24)	6.6 (34)	13.1 (22)	880	14.1 (5)	13.6 (8)	1,387	12.5 (22)
Placer	407	10.3 (49)	8.4 (11)	1,787	7.0 (62)	7.7 (42)	10.8 (11)	595	16.0 (21)	13.6 (8)	1,117	15.2 (47)
Plumas	74	* *	* *	348	4.9 (30)	9.5 (55)	10.3 (9)	139	15.8 (16)	10.8 (2)	180	7.2 (3)
Riverside	6,538	5.9 (15)	7.8 (5)	39,015	5.1 (38)	4.9 (18)	16.3 (51)	15,376	15.9 (18)	15.2 (21)	22,412	14.6 (44)
Sacramento	2,251	6.4 (23)	9.6 (25)	25,229	5.1 (38)	6.8 (37)	13.7 (29)	9,968	15.2 (13)	14.3 (14)	13,685	16.1 (51)
San Benito	134	4.5 (7)	14.2 (52)	1,140	4.2 (19)	5.1 (19)	17.6 (60)	402	19.9 (56)	25.4 (58)	761	14.2 (38)
San Bernar	7,701	7.6 (44)	8.5 (13)	51,757	6.0 (55)	5.7 (25)	15.3 (44)	21,718	15.6 (15)	14.2 (13)	29,015	17.4 (58)
San Diego	8,162	5.8 (13)	10.4 (34)	50,706	4.6 (24)	6.7 (36)	13.3 (25)	19,202	16.0 (21)	14.5 (17)	30,512	11.2 (18)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
San Franci	442	11.1 (51)	7.9 (7)	7,079	5.9 (51)	5.5 (23)	14.9 (40)	2,539	17.5 (46)	19.0 (55)	1,160	9.2 (6)
San Joaqui	3,340	7.1 (37)	8.8 (16)	16,142	6.6 (60)	5.9 (26)	16.1 (50)	6,080	16.7 (37)	18.5 (52)	7,738	10.6 (13)
San Luis O	411	5.8 (13)	11.7 (44)	3,393	3.9 (10)	7.9 (43)	9.8 (6)	1,168	15.8 (16)	12.5 (7)	1,520	12.2 (20)
San Mateo	859	5.9 (15)	10.9 (38)	7,622	5.0 (35)	6.4 (32)	14.2 (35)	2,636	18.9 (52)	19.6 (57)	3,481	7.0 (2)
Santa Barb	2,501	9.5 (48)	6.8 (2)	10,645	4.5 (21)	6.4 (32)	13.6 (28)	3,881	16.5 (34)	17.3 (43)	7,304	11.0 (16)
Santa Clar	1,710	6.1 (20)	9.3 (22)	18,106	4.9 (30)	6.6 (34)	12.9 (20)	5,859	16.1 (24)	15.3 (22)	7,696	14.1 (36)
Santa Cruz	739	7.2 (39)	12.3 (46)	7,484	4.1 (18)	5.1 (19)	14.4 (37)	2,166	16.4 (31)	16.9 (39)	3,648	12.8 (27)
Shasta	754	7.0 (34)	10.9 (38)	3,474	8.2 (64)	4.1 (4)	16.9 (53)	1,386	19.0 (54)	18.2 (51)	2,077	12.5 (22)
Sierra	5	* *	* *	47	* *	* *	* *	20	* *	* *	24	* *
Siskiyou	204	11.8 (52)	7.8 (5)	998	6.0 (55)	7.9 (43)	10.7 (10)	389	16.2 (27)	13.9 (10)	639	20.2 (60)
Solano	969	7.2 (39)	8.8 (16)	5,061	5.8 (48)	8.0 (45)	12.9 (20)	1,631	17.4 (45)	15.6 (25)	2,338	14.5 (43)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Sonoma	879	6.8 (29)	11.4 (42)	4,844	3.7 (8)	9.0 (51)	11.6 (16)	1,600	13.4 (3)	17.3 (43)	1,959	14.8 (45)
Stanislaus	2,443	8.4 (46)	9.4 (23)	12,629	3.8 (9)	9.3 (53)	13.1 (22)	4,737	15.0 (11)	16.4 (33)	9,016	17.0 (55)
Sutter	418	3.3 (1)	12.7 (49)	1,852	2.6 (1)	9.5 (55)	10.9 (12)	698	12.0 (1)	13.9 (10)	1,064	14.9 (46)
Tehama	300	6.7 (27)	8.0 (8)	1,790	3.9 (10)	6.8 (37)	14.0 (32)	719	18.8 (51)	14.3 (14)	1,244	21.9 (62)
Trinity	21	* *	* *	164	6.7 (61)	10.4 (59)	15.2 (43)	70	* *	* *	72	* *
Tulare	2,021	5.2 (8)	10.1 (29)	15,823	5.9 (51)	4.1 (4)	17.5 (59)	5,774	16.7 (37)	16.3 (32)	10,238	14.4 (42)
Tuolumne	60	* *	* *	715	5.9 (51)	6.3 (29)	11.5 (14)	279	14.7 (9)	15.4 (23)	399	13.3 (31)
Ventura	2,766	5.7 (12)	9.8 (27)	12,718	3.9 (10)	5.6 (24)	14.7 (39)	4,454	16.1 (24)	17.0 (40)	8,215	9.5 (9)
Yolo	585	4.4 (6)	12.3 (46)	3,370	3.3 (4)	8.4 (48)	11.7 (18)	1,320	15.9 (18)	15.5 (24)	1,889	14.2 (38)
Yuba	239	7.5 (43)	11.3 (41)	1,338	4.0 (16)	8.8 (50)	12.0 (19)	502	14.3 (7)	13.9 (10)	701	14.1 (36)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this county's rate to other counties. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
California	131,493	6.6	9.1	781,922	5.0	5.3	15.9	306,084	16.3	16.5	445,793	13.7
Nation (Prior Year)	1,172,312	8.9	8.3	4,637,556	6.0	5.4	12.8	1,940,253	15.2	13.2	3,232,431	13.7

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this county's rate to other counties. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alameda	11,503	8.1 (48)	3.2 (53)	18.1 (39)	18.4 (16)	7,703	14.0 (42)
Alpine	70	* *	* *	* *	* *	65	* *
Amador	317	9.8 (60)	1.9 (15)	19.6 (55)	21.1 (46)	277	10.5 (11)
Butte	3,078	8.1 (48)	3.2 (53)	16.0 (8)	18.1 (14)	2,996	11.8 (24)
Calaveras	714	6.7 (22)	1.3 (4)	16.0 (8)	21.0 (44)	697	16.1 (51)
City Of Be	125	4.8 (3)	2.4 (28)	14.4 (5)	27.2 (63)	104	24.0 (61)
Colusa	768	8.6 (54)	1.4 (6)	17.1 (21)	21.5 (52)	514	11.3 (19)
Contra Cos	4,112	7.4 (37)	2.3 (27)	19.1 (51)	21.0 (44)	2,766	15.1 (47)
Del Norte	378	5.0 (5)	0.8 (2)	19.8 (56)	18.5 (19)	361	15.5 (48)
El Dorado	924	6.6 (21)	3.6 (60)	16.5 (16)	15.0 (3)	791	12.4 (30)
Fresno	24,904	8.1 (48)	2.7 (42)	16.4 (11)	19.9 (32)	19,874	11.8 (24)
Glenn	452	8.6 (54)	0.9 (3)	16.4 (11)	24.3 (61)	445	11.2 (17)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Humboldt	1,366	5.3 (10)	1.5 (9)	17.6 (26)	21.2 (48)	1,112	17.5 (55)
Imperial	2,599	5.3 (10)	2.4 (28)	18.0 (37)	23.0 (58)	2,364	8.0 (4)
Inyo	269	7.1 (30)	2.2 (23)	23.8 (63)	21.6 (53)	183	14.2 (44)
Kern	11,093	6.2 (17)	2.6 (38)	17.2 (22)	20.7 (41)	8,915	12.6 (31)
Kings	1,353	4.3 (1)	2.6 (38)	18.8 (48)	20.5 (39)	1,275	16.2 (52)
L.A. East	39,146	7.2 (33)	2.7 (42)	18.5 (44)	20.2 (37)	36,948	11.6 (22)
L.A. North	34,601	7.0 (28)	2.6 (38)	18.2 (40)	18.9 (21)	33,427	9.9 (8)
L.A. Other	47,842	4.4 (2)	2.4 (28)	17.9 (32)	22.8 (57)	35,397	12.6 (31)
L.A. South	55,386	8.1 (48)	2.5 (35)	18.6 (46)	18.6 (20)	54,121	10.7 (14)
L.A. West	12,678	5.7 (12)	2.4 (28)	18.0 (37)	19.7 (28)	11,126	14.1 (43)
Lake	524	7.1 (30)	1.5 (9)	20.0 (58)	19.5 (25)	455	13.2 (38)
Lassen	251	5.2 (8)	2.4 (28)	16.3 (10)	15.9 (5)	201	7.0 (3)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Long Beach	6,636	5.1 (6)	2.4 (28)	17.8 (29)	21.4 (50)	5,283	14.4 (46)
Madera	2,748	7.6 (40)	2.7 (42)	17.4 (25)	20.1 (36)	3,026	13.3 (40)
Marin	783	11.0 (63)	1.9 (15)	20.4 (61)	21.7 (54)	550	6.9 (2)
Mariposa	100	10.0 (62)	3.0 (48)	14.0 (4)	14.0 (2)	96	* *
Mendocino	1,048	7.5 (38)	1.7 (11)	19.9 (57)	19.6 (27)	671	12.1 (29)
Merced	4,094	9.1 (58)	2.0 (19)	18.9 (49)	19.7 (28)	4,216	19.7 (59)
Modoc	141	9.9 (61)	0.0 (1)	13.5 (2)	25.5 (62)	132	9.8 (7)
Mono	110	7.3 (36)	3.6 (60)	16.4 (11)	12.7 (1)	104	21.2 (60)
Monterey	3,091	7.9 (44)	1.9 (15)	19.3 (53)	21.1 (46)	3,301	11.5 (21)
Napa	805	8.2 (52)	2.6 (38)	18.5 (44)	20.0 (35)	341	18.8 (57)
Nevada	361	7.2 (33)	3.3 (58)	16.6 (17)	15.5 (4)	255	6.3 (1)
Orange	40,731	6.8 (23)	3.1 (51)	17.9 (32)	19.4 (24)	35,106	13.1 (37)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Pasadena	1,215	5.1 (6)	2.9 (47)	17.9 (32)	18.4 (16)	1,104	10.5 (11)
Placer	817	8.9 (57)	5.1 (63)	17.6 (26)	17.3 (7)	634	9.3 (6)
Plumas	199	6.0 (14)	2.5 (35)	17.6 (26)	16.6 (6)	158	10.8 (16)
Riverside	24,039	6.2 (17)	3.0 (48)	17.2 (22)	19.7 (28)	21,756	13.4 (41)
Sacramento	16,658	7.5 (38)	3.2 (53)	16.4 (11)	17.5 (10)	11,855	16.0 (50)
San Benito	603	7.1 (30)	1.3 (4)	19.1 (51)	23.1 (59)	559	10.7 (14)
San Bernar	33,176	6.0 (14)	3.2 (53)	16.9 (18)	19.0 (22)	28,180	14.3 (45)
San Diego	22,649	6.3 (19)	2.5 (35)	17.9 (32)	19.5 (25)	20,497	12.6 (31)
San Franci	5,052	8.0 (47)	3.1 (51)	15.7 (6)	18.1 (14)	1,939	11.9 (28)
San Joaqui	6,889	8.8 (56)	3.0 (48)	16.4 (11)	19.9 (32)	4,381	11.3 (19)
San Luis O	1,295	7.6 (40)	2.4 (28)	17.9 (32)	18.4 (16)	772	11.7 (23)
San Mateo	3,783	9.1 (58)	1.7 (11)	20.6 (62)	22.4 (56)	2,695	8.9 (5)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Santa Barb	3,513	7.9 (44)	2.0 (19)	20.2 (60)	22.0 (55)	3,459	10.2 (10)
Santa Clar	7,631	7.9 (44)	3.3 (58)	17.0 (20)	20.8 (42)	4,615	12.8 (35)
Santa Cruz	2,662	6.0 (14)	1.7 (11)	20.0 (58)	23.6 (60)	2,295	12.7 (34)
Shasta	1,428	6.9 (26)	1.4 (6)	16.9 (18)	17.9 (13)	1,087	13.2 (38)
Sierra	29	* *	* *	* *	* *	27	* *
Siskiyou	574	5.2 (8)	2.1 (21)	18.3 (42)	19.9 (32)	531	16.8 (54)
Solano	2,312	6.5 (20)	2.2 (23)	17.8 (29)	20.6 (40)	1,828	19.3 (58)
Sonoma	2,062	7.6 (40)	1.4 (6)	18.6 (46)	20.3 (38)	1,243	11.2 (17)
Stanislaus	5,531	5.9 (13)	2.8 (46)	18.2 (40)	19.1 (23)	5,680	15.5 (48)
Sutter	661	6.8 (23)	4.8 (62)	15.7 (6)	17.5 (10)	522	17.8 (56)
Tehama	908	7.0 (28)	1.9 (15)	19.4 (54)	17.3 (7)	903	16.4 (53)
Trinity	117	8.5 (53)	1.7 (11)	11.1 (1)	19.7 (28)	75	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2001 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Comparison of Growth and Anemia Indicators by County (3,4)							
County	Height and Weight					Anemia Low Hb/Hct (7)	
	Short Stature (5)	Under- weight (6)	Over- weight (6)				
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Tulare	7,950	7.2 (33)	2.2 (23)	18.3 (42)	21.2 (48)	7,406	13.0 (36)
Tuolumne	318	6.9 (26)	2.2 (23)	18.9 (49)	17.3 (7)	238	10.5 (11)
Ventura	5,101	4.8 (3)	3.2 (53)	17.2 (22)	20.9 (43)	5,143	10.0 (9)
Yolo	1,552	6.8 (23)	2.1 (21)	17.8 (29)	21.4 (50)	1,227	11.8 (24)
Yuba	668	7.6 (40)	2.7 (42)	13.5 (2)	17.8 (12)	626	11.8 (24)
California	474,493	6.8	2.7	17.8	19.9	406,633	12.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.