


QUICK GUIDE: MAT Use for Opioid Use Disorder



| | BUPRENORPHINE | METHADONE | NALTREXONE |
|---------------------------|---|--|--|
| COMMON BRANDS | Suboxone, Zubsolv, Bunavail, Subutex, Probuphine, Sublocade | Methadose, Diskets, Dolophine | Vivitrol |
| TYPE | | | |
| HOW IT WORKS | <ul style="list-style-type: none"> • Makes the brain think it is still getting the problem opioid. Prevents cravings and withdrawal symptoms and reduces the risk of overdose. • Buprenorphine can be prescribed by a trained provider in a doctor's office or other health care setting, as well as in a narcotic treatment program (NTP). | <ul style="list-style-type: none"> • Makes the brain think it is still getting the problem opioid. Prevents cravings and withdrawal symptoms and reduces the risk of overdose. • Methadone is dispensed only in highly regulated NTPs. | <ul style="list-style-type: none"> • Blocks the effects of opioids. • Naltrexone is not a controlled substance and can be prescribed or administered in any health care or substance use disorder (SUD) setting, such as a doctor's office or clinic. |
| THINGS TO CONSIDER | <ul style="list-style-type: none"> • Treatment can start quickly, as soon as someone enters withdrawal. • Flexible dosing schedule. • Relapse risk increases if you forget or choose not to take medication. • Common side effects are headache, nausea, and constipation. | <ul style="list-style-type: none"> • Treatment can start right away, no need for detoxification. • Less flexible schedule. Dosing occurs in the early morning. • Side effects include constipation, sexual problems, swelling, and sweating and potential heart problems. | <ul style="list-style-type: none"> • Less evidence for effectiveness in OUD treatment than buprenorphine or methadone. • Does not cause physical dependence. • Not recommended for pregnant women as detox can harm the baby. Methadone or buprenorphine are recommended for pregnant women with OUD. |

CONTINUED ON BACK

QUICK GUIDE: MAT Use for Opioid Use Disorder

| CONTINUED FROM FRONT | BUPRENORPHINE | METHADONE | NALTREXONE |
|------------------------------|---|--|--|
| THINGS TO CONSIDER | <ul style="list-style-type: none"> • Causes physical dependence. If or when you want to come off the drug, you will need to do so slowly to minimize the discomfort of detox symptoms. • Buprenorphine is sometimes used short term to relieve pain associated with detox, but more often used long term, known as maintenance treatment. | <ul style="list-style-type: none"> • Causes physical dependence. If you want to come off the drug, you will need to do so slowly to minimize the discomfort of detox symptoms. • May cause drowsiness at first before maintenance dose is determined. • Methadone is often a good option for people who have used opioids for a long time or have been unsuccessful with other treatments. | <ul style="list-style-type: none"> • Detox from opioids is required before taking naltrexone. Side effects may include stomach pain, nausea, vomiting, headache, joint pain, trouble sleeping and anxiety. Some people also report soreness in the area of the injection. • Injection form of the medication lasts for about 30 days before it wears off. Overdose risk can be higher after naltrexone wears off due to decrease in tolerance. |
| QUESTIONS FOR CLIENTS | <ul style="list-style-type: none"> • Can you commit to taking this medication daily? • Are you comfortable with taking a medication that requires time to taper off to minimize the discomfort of detox? | <ul style="list-style-type: none"> • Have you used opioids for a long time, or have you found other treatments have not worked well for you? • Can you come to the clinic in the early morning for dosing? Will you need to make arrangements for work or transportation? • Do you work in an industry with heavy machinery? Could your work be affected by possible drowsiness during your initial dosing period? • Are you comfortable with taking a medication that requires time to taper off to minimize the discomfort of detox? | <ul style="list-style-type: none"> • Have you detoxed from opioids, or would you be willing to detox to take this medication? • Can you commit to making an appointment once every month to continue receiving the injection? • Do you have any medical needs that would be affected by blocking the opioid receptors? For example, do you use opioids to reduce chronic pain? |

| | NALTREXONE | ACAMPROSATE | DISULFIRAM |
|---------------------------|--|---|--|
| COMMON BRANDS | Revia, Vivitrol | Campral | Antabuse |
| TYPE |  |  |  |
| HOW IT WORKS | <ul style="list-style-type: none"> • Medication that blocks the effects of alcohol and reduces cravings. • Offered as a daily pill or monthly injection. • Naltrexone is not a controlled substance and can be prescribed or administered in any health care or SUD setting, such as a doctor's office or clinic. | <ul style="list-style-type: none"> • Medication to reduce cravings for patients who have already stopped drinking. It does not help with withdrawal symptoms but does reduce cravings. If relapse occurs, patients can continue taking the medication without needing to detox first. • Offered as a tablet taken three times a day. • Acamprosate is not a controlled substance and can be prescribed or administered in any health care or SUD setting, such as a doctor's office or clinic. | <ul style="list-style-type: none"> • Medication that causes severe vomiting if someone drinks alcohol. • Offered as daily pill. • Disulfiram is not a controlled substance and can be prescribed or administered in any health care or SUD setting such as a doctor's office or clinic. |
| THINGS TO CONSIDER | <ul style="list-style-type: none"> • Detoxification from alcohol is required before taking naltrexone. • Relapse risk increases if you forget or choose not to take pill form of the medication. | <ul style="list-style-type: none"> • Detoxification from alcohol is not required but is highly recommended before starting on acamprosate. • Relapse risk increases if patients forget or choose not to take medication. | <ul style="list-style-type: none"> • Detoxification from alcohol is required. • Relapse risk increases if you forget or choose not to take medication. |

QUICK GUIDE: MAT Use for Alcohol Use Disorder

| CONTINUED FROM FRONT | NALTREXONE | ACAMPROSATE | DISULFIRAM |
|------------------------------|---|---|--|
| THINGS TO CONSIDER | <ul style="list-style-type: none">• Injection form of the medication lasts for about 30 days before it wears off. | <ul style="list-style-type: none">• Common side effects include stomach pain, dizziness or dry mouth; more rarely patients may experience anxiety or depression. | <ul style="list-style-type: none">• Side effects are not common but may include headache, drowsiness or rash.• Disulfiram can be a good option for compulsive drinking (everything is fine and then you have a strong urge to drink). |
| QUESTIONS FOR CLIENTS | <ul style="list-style-type: none">• Have you detoxed from alcohol, or would you be willing to detox to take this medication?• Can you commit to taking this medication daily, or would a month-long injection be a better option?• Do you have any medical needs that would be affected by blocking the opioid receptors? For example, do you use opioids to reduce chronic pain? | <ul style="list-style-type: none">• Can you commit to taking this medication three times a day?• Do you feel that craving reduction alone is enough to help you stop drinking, or do you need something more? For example, disulfiram makes you vomit if you drink, and naltrexone takes away the pleasurable feeling of drinking. | <ul style="list-style-type: none">• Have you detoxed from alcohol, or would you be willing to detox to take this medication?• Can you commit to taking this pill daily?• Do you work in an industry with exposure to alcohol-based products (i.e., paint thinner, varnish, etc.) which could react with the medication?• Are you willing to run the risk of severe vomiting should you relapse? |